

Global Alliance for the Promotion of Physical Activity: the Hamburg Declaration

Jürgen M Steinacker ^{1,2,3,4} Willem van Mechelen,^{2,5,6,7,8} Wilhelm Bloch,^{9,10} Mats Börjesson,^{2,11,12} Maurizio Casasco,¹³ Bernd Wolfarth,^{3,14,15} Carolin Knoke ^{1,2} Theodora Papadopoulou,^{16,17} Janine Wendt,¹ Hashel Al Tunajji,^{18,19} Dietrich Andresen,²⁰ Olena Andrieieva,²¹ Norbert Bachl,^{22,23} Victoriya Badtieva,^{24,25} Friedhelm J Beucher,²⁶ Cheri A Blauwet,²⁷ Jose-Antonio Casajus Mallen ^{28,29,30} Ju-Ho Chang,³¹ German Clénin,^{32,33} Naama Constantini,^{34,35} Demitri Constantinou ^{36,37} Luigi Di Luigi,³⁸ Lukas Declercq,³⁹ Stephane Doutreleau ^{40,41} Svitlana Drozdovska,⁴² Martine Duclos ^{41,43,44} Andrea Ermolao,^{45,46} Thomas Fischbach,⁴⁷ Anastasia N Fischer,^{48,49} Chiara Fossati,⁵⁰ George Franchella,⁵¹ Mark Fulcher,^{52,53} Jan C Galle,⁵⁴ Christian Gerloff,⁵⁵ Evelina Georgiades,⁵⁶ Boris Gojanovic,^{57,58} Marcela González Gross,^{30,59} Andy Grote,⁶⁰ Martin Halle,^{61,62} Hans Hauner,⁶³ Matthew Payton Herring ⁶⁴ Mikio Hiura,⁶⁵ Kerstin Holze,⁶⁶ Gerhard Huber,^{67,68} David Hughes,^{69,70} Mark R. Hutchinson,^{49,71,72} Anca Ionescu,^{73,74} Dina Christina Janse van Rensburg ^{37,75} Anna Jegier,^{73,76} Natasha Jones,⁷⁷ Kirsten Kappert-Gonther,⁷⁸ Monika Kellerer,⁶³ Yutaka Kimura,^{79,80} Agrita Kiopa,⁸¹ Bernd Kladny,⁸² Gerhard Koch,⁸³ Elin Kolle,⁸⁴ Greg Kolt,⁸⁵ Yiannis Koutedakis,^{86,87} Stephan Kress,⁸⁸ Susi Kriemler ^{33,89} Jens Kröger,⁹⁰ Christian Kuhn,^{91,92} Roman Laszlo,⁹³ Ralph Lehnert,⁹⁴ François J Lhuissier ^{41,95,96} Kerstin Lüdtke,⁹⁷ Shigeru Makita,^{80,98} Pedro Manonelles Marqueta,^{23,98} Winfried März,⁹⁹ Kirill Micallef-Stafrace,^{73,100} Mike Miller,¹⁰¹ Melita Moore,¹⁰² Erich Müller,¹⁰³ Daniel Neunhäuserer,^{45,46} I. Renay Onur,¹⁰⁴ Vahur Ööpik,¹⁰⁵ Malgorzata Perl,¹⁰⁶ Anastassios Philippou,⁸⁶ Hans-Georg Predel,^{107,108} Sebastien Racinais,¹⁰⁹ Algirdas Raslanas,¹¹⁰ Ruediger Reer,^{2,73,111} Klaus Reinhardt,¹¹² Claus Reinsberger,¹⁵ Sandra Rozenstoka,^{23,81,113,114} Robert Sallis ¹¹⁵ Luis B Sardinha ^{116,117} Martin Scherer,^{118,119} Jasper Schipperijn,¹²⁰ Romain Seil,¹²¹ Benedict Tan ^{122,123} Arno Schmidt-Trucksäss,¹²⁴ Nils Schumacher,¹¹¹ Bernhard Schwaab,¹²⁵ Ansgar Schwartz,¹²⁶ Masato Suzuki,⁸⁰ Jeroen Swart,^{23,127} Ralph Tiesler,¹²⁸ Ulf Tippelt,¹²⁹ Eleanor Tillet,^{17,130} Jane Thornton ¹³¹ Bulent Ulkar ^{23,132} Eve Unt,¹³³ Evert Verhagen ⁵ Thomas Weikert,⁶⁶ Roberto Vettor,^{46,134} Sheng Zeng,^{23,135} Richard Budgett,¹³⁶ Lars Engebretsen,^{136,137} Ugur Erdener,¹³⁶ Fabio Pigozzi,^{3,138} Yannis P Pitsiladis ^{3,139}

To cite: Steinacker JM, Mechelen W, Bloch W, et al. Global Alliance for the Promotion of Physical Activity: the Hamburg Declaration. *BMJ Open Sport & Exercise Medicine* 2023;**9**:e001626. doi:10.1136/bmjsem-2023-001626

Accepted 29 June 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Jürgen M Steinacker;
juergen.steinacker@uni-ulm.de

ABSTRACT

Non-communicable diseases (NCDs), including coronary heart disease, stroke, hypertension, type 2 diabetes, dementia, depression and cancers, are on the rise worldwide and are often associated with a lack of physical activity (PA). Globally, the levels of PA among individuals are below WHO recommendations. A lack of PA can increase morbidity and mortality, worsen the quality of life and increase the economic burden on individuals and society. In response to this trend, numerous organisations came together under one umbrella in Hamburg, Germany, in April 2021 and signed the 'Hamburg Declaration'. This represented an international commitment to take all necessary

actions to increase PA and improve the health of individuals to entire communities. Individuals and organisations are working together as the 'Global Alliance for the Promotion of Physical Activity' to drive long-term individual and population-wide behaviour change by collaborating with all stakeholders in the community: active hospitals, physical activity specialists, community services and healthcare providers, all achieving sustainable health goals for their patients/clients. The 'Hamburg Declaration' calls on national and international policymakers to take concrete action to promote daily PA and exercise at a population level and in healthcare settings.

Box 1 Definitions of physical activity, physical inactivity, sedentary behaviour as well as guidelines and recommendations on physical activity provided by the World Health Organization (WHO)

Physical activity (PA) can be defined as any bodily movement produced by skeletal muscles that requires energy expenditure.² Physical inactivity⁵ refers to any waking behaviour while in a sitting, reclining or lying posture with low energy expenditure. Sedentary behaviour is any waking behavior characterized by an energy expenditure ≤ 1.5 metabolic equivalents (METs), while in a sitting, reclining or lying posture.⁶ Passive standing is an intermediate activity with ≤ 2.0 METs.⁶ How much PA - depending on age groups and population groups - is required for a good health is provided in the guidelines and recommendations of the World Health Organization (WHO). For example, adults aged 18 to 64 years should do at least 150 minutes of moderate-intensity aerobic PA or at least 75 minutes of intense aerobic PA throughout the week.^{6,28}

BENEFITS OF PHYSICAL ACTIVITY

Low levels of physical activity (PA) in the population and sedentary lifestyle^{1,2} (see definitions in [Box 1](#)) have been associated with the global rise in non-communicable diseases (NCDs), including coronary heart disease, stroke, hypertension, type 2 diabetes, dementia, depression and certain cancers (eg, bladder, breast, colon) and musculoskeletal disorders.^{3,4} Moreover, physical inactivity is the fourth leading risk factor for premature mortality.² The WHO stipulates the amount of PA needed for good health (see [Box 1](#)).^{5,6} The high prevalence of insufficient PA across the globe from 2001 to 2016 can be interpreted as worrying ([figure 1A](#) for women and [figure 1B](#) for men).⁷

Regular exercise, in particular aerobic exercise such as walking, hiking or cycling,⁸ is recommended for the prevention, therapy and rehabilitation of many NCDs, such as cardiovascular,⁹ metabolic,¹⁰ neurodegenerative diseases^{11,12} and specific types of cancer,¹³ in addition to their important role in the management of depression and anxiety disorders.¹⁴ Personalised PA prescriptions, which include variations in the type of exercise, intensity, frequency and duration, have many positive health benefits - not only confined to physical but also to psychosocial function,¹⁵ well-being and quality of life.¹⁶ PA prescription may provide equal or even better therapeutic outcomes for various health conditions than other standard medical treatments.¹⁷ For specific musculoskeletal injuries, PA prescription is known to be an essential part of therapy.³ As little as 15 min a day of moderate-intensity PA can help lower the risk for cardiovascular disease.¹⁸ However, the effects of an individually tailored programme are even more effective.^{4,19,20}

Although there is respective guidance to promote PA as part of standard healthcare, there is a lack of a concerted effort to increase PA as a preventive health measure and to maintain quality of life, which could reduce the burden

of NCDs and associated risk of mortality and morbidity, as well as direct and indirect costs to the healthcare system.⁴ This is especially true in NCDs such as obesity,^{8,21,22} type 2 diabetes,²³ heart disease,²⁴ hypertension,²⁵ cancer²⁶ and stroke.²⁷ The 'Step up! Tackling the burden of insufficient PA in Europe' report from the WHO and the Organisation for Economic Co-operation and Development (OECD) estimates that every euro invested in PA would save around €1.7 in NCDs health expenses²⁸ and the global cost of inaction on physical inactivity was calculated to reach approximately €44.3 billion per year, up to 2030.²⁹

The WHO recommends various evidence-based policy actions to create active societies, environments, people and systems.³⁰ Such a comprehensive whole-system approach,³⁰ and ecological and multilevel actions that consider personal, environmental and political factors, are needed to address physical inactivity.³¹ So far, such approaches have already been used to reduce tobacco consumption³² and to create food environments supporting healthy eating behaviours.³³ Such approaches should also be increasingly considered and used when creating an environment conducive to PA. Many countries worldwide have already developed formal written policies, guidelines or targets to promote PA.³⁴ However, knowledge about the implementation and effectiveness of PA policies is still limited.³⁵ Moreover, no clear guidance exists for governments on which policies are preferable in different settings and under various preconditions.³⁶⁻³⁸ In response, the 'Global Alliance for the Promotion of Physical Activity' members propose an approach that addresses both the promotion of PA in society and is tailored to the individual. This requires an integrated health policy that takes into account the need to increase PA²² at the population level³⁹ as well to integrate individual PA prescriptions as a standard of care.⁴⁰

Scientific societies and organisations working towards global health through PA

Despite the overwhelming evidence of the benefits of PA and exercise for general health prevention and treatment of NCDs, it remains an auxiliary therapy in most countries




Figure 1 Country prevalence of insufficient physical activity (A) in women in 2016 and (B) in men in 2016.⁷

rather than the first choice in standard treatment and care. In response to this impasse, 139 organisations from different countries have indicated their full commitment by signing the ‘Hamburg Declaration’. The so-called Global Alliance (GA) has agreed to jointly pursue the goal of promoting PA as a primary preventive measure to improve and maintain the health of populations worldwide and to facilitate its integration into daily patient care. The GA aims to encourage and support national and international policymakers, health professionals and providers, and other health sector actors to introduce and implement the actions required at various levels (political, socio-political and social, [figure 2](#)).

Joining global efforts

To achieve this, the GA plans several actions based on evidence-based practices from around the world,^{4 6 38 41} drawing on current research findings as well as marketing,⁴² communication and dissemination strategies.⁴³ Overall, the GA attaches particular attention to close cooperation between all members and the engagement of setting-specific actors (eg, the general population, community-based healthcare, hospitals, research organisations and politicians). Such an engagement of various actors is an ongoing process of soliciting knowledge, experience, judgement and values from the people involved,⁴⁴ including the target population.⁴⁵ It also serves to create a shared understanding, make transparent decisions⁴⁶ and understand the interests of various parties.^{37 47}

There are different tools to achieve successful participation,⁴⁸ and engaging various actors is crucial to granting shared decision-making capacity to those who traditionally have limited authority, ensuring an equitable dynamic with powerful actors. A synergistic and collective approach will lead to outcomes and elicit change that no one constituent member could have produced on their own - this is the foundation for future activities of the GA. This means the need to introduce behavioural techniques like the ‘Theory of Change’ methodology for interacting with actors⁴⁸ or shared decision-making with patients.⁴⁹

The synergies created between the GA members and the respective actors will ensure a dynamic and effective fight against the worldwide pandemic of PA,⁵⁰ including initiating change at the highest level (eg, policies determined at the governmental level). Each actor fulfils different and synergistic roles and objectives, contributing to the overall goal. For example, the Organising Committee of the Paris 2024 Summer Olympics is using the energy and strong reputation of the Games to promote the ‘30 min of PA per day at school’ message. The IOC and the WHO signed a memorandum to increase levels of daily PA among the world’s population, to support NCD prevention and healthier lifestyles through sport and exercise, and to enhance cooperation between the health and the sports sector.⁵¹ The first initiative was planned for Olympic Day, 23 June 2023, with the handle ‘#LetsMove’




Figure 2 Population-based health promotion and the interaction of various actors at different levels to support and communicate with the target group in order to achieve health goals.

and the message ‘Move your body and dedicate 30 min to get active with Olympians’.⁵² The International Federation of Sports Medicine (FIMS) supports the promotion of a healthy and active lifestyle through high-quality education of (para-) medical professionals and the implementation of evidence-based sports and exercise medicine.^{51 53}

Call for including exercise and sport in health policies

Studies have been demonstrating that population health is influenced by the environment in the broadest sense,^{37 38 54} including economic factors,⁵⁵ access to green spaces, water and air quality,^{54 56} employment, access to medical care,⁵⁷ the availability of safe non-motorised transportation (eg, walking, cycling) systems,^{30 54 58} as well as education.^{35 55 59} In this framework, individual behaviour reflects individuals’ personal responsibility through health-promoting behaviours and choices - of which PA is one of the most important.^{13 21 59 60} Heritability affects the likelihood of developing a particular NCD.^{8 26 61} However, lifestyle, particularly PA, can modify genetically determined expression patterns through epigenetic changes towards reducing the frequency or severity of NCD and, thus, a better quality of life.^{4 13 62–64}

The constantly high prevalence of physical inactivity from 2001 to 2016 suggests that the target set by WHO Member States to reduce physical inactivity by 10% by 2025 will not be achieved, despite it is as one of nine WHO targets for preventing and treating NCDs.³⁰ The forced lockdown during the COVID-19 pandemic contributed further to the rise in physical inactivity,³⁸ and physical inactivity per se is considered an important independent risk factor for severe sequelae and death.^{31 65} Global PA has only partially been restored¹⁴ and remains an enduring negative ‘side effect’ of the COVID-19 pandemic.⁶⁶ The

reasons behind this are multi-fold⁶⁷; lack of awareness by the general population of the health benefits of PA,³⁵ lack of policy definition and efforts, shortfalls in the education and training provided to medical practitioners on the subject, lack of specialised practitioners trained in prescribing and monitoring exercise programmes. The need, therefore, for a concerted, dynamic action that advocates the inclusion of PA and exercise in health policies has never been greater.

The critical mass created by the concerted efforts of the GA is expected to enable national and global policymakers to prioritise promoting PA and exercise as medicine and supporting population health. A systematic global interaction is needed among all participants, including individuals, organisations and communities with a direct interest in the process and outcomes of projects, research or policy.^{38 44} This work should target the sport, school, health and medicine sectors and focus on successfully implementing PA and exercise promotion measures.^{30 59} Policies should create an ecosystem that promotes healthy lifestyles, reduces sedentary behaviour and integrates PA and exercise as medicine in the daily care of patients.^{37 55 68 69} For example, the WHO has produced a series of guides and toolkits to help implement best practices for integrating PA into the daily lives of the general population and into patient care and rehabilitation.^{56–58} Collaboration among organisations led to an effort by scientists from all fields to work on this common cause, as reflected in a large number of multi-disciplinary publications.

In most countries, sports and exercise science/medicine is not part of the general education programme for students.⁷⁰ Sports and exercise medicine specialists should be the best qualified to prescribe PA for disease prevention and treatment. However, in terms of specialisation, sports and exercise science/medicine is not part of the training of medical specialists in most countries.⁷¹ These deficiencies should be addressed, and the goal should be for every medical professional to be able to use PA as a therapy in their respective clinical settings.

In most countries, recommendations for PA during hospitalisation and at discharge are not mentioned, and PA and exercise in the hospital and at home are extremely rare in treatment.⁴ The WHO has already called for the introduction and strengthening of patient assessment and counselling systems to increase PA and reduce physical inactivity in the general population⁶ through appropriately trained health, community and social service providers in primary and secondary care and social services as part of general healthcare.^{30 57}

The role of active sport cities

When health professionals focus traditionally mainly on the individual, in future, medical stakeholders have to include a population approach. Otherwise, large parts of the community may lose out on benefitting.^{39 72} Population-based strategies to promote PA should include a healthy ecosystem and the resources necessary to lead an active life

and integration into the healthcare system.⁶⁹ Active societies can achieve more than the simple provision of health benefits. For example, an active society conserves health system resources, cleans the air and ensures less congestion and safer streets.^{38 56} In addition, opening schoolyard PA facilities during public holidays is an effective way to promote PA among children in a neighbourhood. We call for and support the concept of ‘global active and smart sports cities’ with active, smart and sustainable hospitals where citizens, stakeholders, organisations and policymakers, working together, can be models for linking the healthcare system to public health, integrating PA and exercise as medicine.^{60 69} The ‘global active and smart sports city’ concept, conceived in Hamburg in April 2021, builds on the global active city idea, and this new concept is now being developed for major cities part of the GA, including Olympic cities that will address preventive measures by community interaction.^{60 73} The concept aims to integrate PA into everyday urban life and to create the conditions to enable citizens to live active lives through interaction with smart technology and urban planning to improve visual attractiveness, create a more sustainable transportation system and promote active commuting (eg, walkability and bicycle friendliness).^{56 58 60 68}

Active hospitals achieve sustainable health goals for their patients

Healthcare should be more sustainable, and so-called ‘active hospitals’ that incorporate PA into patients’ treatment plans could be a model of the future. The goal is to systematically integrate PA and exercise interventions into patient care.^{3 4 9 10 20} This could increase patients’ PA, improve staff communication skills and build collaboration with local sports clubs, sports and fitness specialists, other community services and self-help organisations. The ‘Exercise is Medicine’ initiative,⁴¹ ‘Moving Medicine’,⁷⁴ the Swedish ‘PAP (Physical Activity on Prescription)’ model²⁰ or the Dutch ‘Exercise=Medicine’ model⁷⁵ are excellent examples of PA prescription initiatives.

Personalised prescriptions and the role of technology

One reason physicians do not offer PA prescriptions in their practices could be a lack of time.⁷⁰ Still, they are often unfamiliar with a patient’s current PA patterns and context factors (eg, environment, family and social support).^{35 70 76} An individualised prescription, including NCD-specific PA and exercise recommendations, could be more beneficial than standard PA prescriptions.^{4 17 19} However, standard PA prescriptions achieve equal or even better therapeutic outcomes than traditional medical treatments.^{17 19 75}

Evidence suggests that the ‘ideal’ PA prescription for a given NCD or risk factor must consider person-related factors⁴⁹ or the individual environment,^{38 76} which may lead to a personalised prescription that includes variations in the type of exercise, intensity, frequency and duration.¹⁹ However, the effects of an individually tailored programme can be even more effective.^{10 29} Precision medicine is driving individualised treatment of NCDs,⁷⁷ and in future, the knowledge of genetic predisposition could further contribute to

prescription.⁷⁸ Through shared decision-making, the patient should decide whether to make positive lifestyle changes, such as increasing PA.⁴⁹

Technology can increase participation and adherence to PA and especially improve the personalisation of PA and its management. Personalisation of interventions using technology is essential, as target users vary widely regarding their PA levels, requirements, preferences and behaviours.⁷⁹ Technology like consumer wearables (eg, smart devices) can influence the general population to the extent that could lead to greater acceptance of PA, more enjoyment and better health outcomes and well-being.⁸⁰

What needs to be changed?

The GA will champion efforts to increase the uptake of PA and exercise for general health and prescription in medical conditions. This will be achieved by:

- ▶ Promoting the inclusion of exercise and sport in health policies.
- ▶ Supporting all measures to increase PA.
- ▶ Facilitating the integration of PA and exercise into daily patient care.

Practical examples include the development of more evidence-based practices, knowledge, experience, judgement and values in the education and training courses for medical students, physicians and other healthcare providers. Particular priority areas will include promoting active and smart sports cities, active hospitals and personalised prescriptions assisted by innovative technologies.

One next step of the GA will be to address the deficits in the education/training and implementation of sports and exercise medicine and to propose a road map for prevention, clinical care and treatment. With this goal in mind, the GA will address which discipline(s) are best suited to prescribe PA exercises and which to administer.

Conclusions

The world is suffering from an obesity epidemic from declining PA. Every age group is affected, especially children. Lack of PA causes so many problems for individuals and society. Diabetes, heart conditions, depression, blood disorders, the unsustainable rising burden and cost on our health services, escalating infrastructure costs and millions in lost productivity due to those unable to work. The solution is to get more people moving. That is the overarching aim of the ‘Global Alliance for the Promotion of Physical Activity’, supported by over 139 organisations worldwide, including the IOC and enshrined in the Hamburg Declaration. Getting people moving, from whatever individual baseline level, as a gateway to greater PA and exercise (and sport) is the critical priority. Working together, we have a better chance to achieve this ambitious goal by creating the initiatives to do this and influencing the decision-makers to invest in PA for the benefit of all. Achieving this goal will save lives, save money, make people happier and help bring the world together. Here are five key messages being championed by the GA:

- ▶ **Promote PA as medicine.** Use influencers and role models in all aspects of life to deliver the message that health, wellness and happiness begin at home. That PA is the best medicine. That prevention is better than cure.
- ▶ **Lobbying decision-makers.** Create the toolkits and systems so that all members of the GA can lobby governments, businesses and non-governmental organisations on at all levels to invest in PA to save lives, save money and improve society.
- ▶ **Adapting PA to the individual, community and their surroundings.** PA must be tailored to the individual, their age, gender, socioeconomic and cultural realities as well as climatic conditions.
- ▶ **Leverage the latest tech.** Work with the big tech, pharma and medical companies and educational establishments to build PA into daily life through wearable devices, smartphones, the internet and the metaverse. Make it easy and fun for people to get into PA through gamification, rewards and community building.
- ▶ **A call for more trials on effectiveness and implementation of policies and programs.** Most information in prevention comes from cohort studies, and there is an urgent need for well-designed trials for physical activity in populations and the healthcare system - particularly in respect of underserved minorities.

Author affiliations

¹Division of Sports and Rehabilitation Medicine, University Hospital Ulm, Ulm, Germany

²European Initiative for Exercise in Medicine (EIEIM), Ulm, Germany

³International Federation of Sports Medicine, Fédération Internationale de Médecine du Sport (FIMS), Lausanne, Switzerland

⁴Institute for Rehabilitation Medicine Research at Ulm University, Institut für rehabilitationsmedizinische Forschung an der Universität Ulm, Bad Buchau, Germany

⁵Department of Public and Occupational Health, location Vrije Universiteit, Amsterdam University Medical Centers, Amsterdam, Netherlands

⁶School of Human Movement and Nutrition Sciences, Faculty of Health and Behavioural Sciences, University of Queensland, Brisbane, Queensland, Australia

⁷Division of Exercise Science and Sports Medicine (ESSM), Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

⁸UCD School of Public Health, Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland

⁹Institute for Cardiology and Sports Medicine, German Sport University, Cologne, Germany

¹⁰Exercise is Medicine Germany, Frankfurt, Germany

¹¹Department of Molecular and Clinical Medicine, University of Gothenburg, Goteborg, Sweden

¹²Institute of Medicine, Sahlgrenska University Hospital, Goteborg, Region Västra Götaland, Sweden

¹³Italian Federation of Sports Medicine (FMSI), Rome, Italy

¹⁴Department of Sport Medicine, Humboldt University and Charité University School of Medicine, Berlin, Deutschland, Germany

¹⁵German Society for Sports Medicine and Prevention, Deutsche Gesellschaft für Sportmedizin und Prävention (DGSP), Frankfurt, Germany

¹⁶Defence Medical Rehabilitation Centre, Stanford Hall, Loughborough, UK

¹⁷British Association of Sport and Exercise Medicine, Doncaster, UK

¹⁸Sports Medicine, United Arab Emirates National Olympic Committee, Dubai, UAE

¹⁹Sports Medicine & Sciences Unit, Zayed Military University, Abu Dhabi, UAE

²⁰German Heart Foundation, Die Deutsche Herzstiftung, Berlin, Germany

- ²¹Department of Health, Fitness and Recreation, National University of Physical Education and Sport of Ukraine, Kiev, Ukraine
- ²²Institute of Sports Science, University of Vienna, Vienna, Austria
- ²³International Federation of Sports Medicine, Lausanne, Switzerland
- ²⁴Sport Medicine, I M Sechenov First Moscow State Medical University, Moscow, Russia
- ²⁵Sport Medicine, Moscow Scientific and Practical Center of Medical Rehabilitation and Sports Medicine, Moscow, Russian
- ²⁶National Paralympic Committee Germany (Deutscher Behindertensportverband (DBS)), Bonn, Germany
- ²⁷Department of Physical Medicine and Rehabilitation, Spaulding Rehabilitation Hospital and Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA
- ²⁸University of Zaragoza, GENUd "Growth, Exercise, Nutrition and Development" Research Group, Zaragoza, Spain
- ²⁹Department of Psychiatry and Nursing, Faculty of Health and Sport Science (FCSD), University of Zaragoza, Zaragoza, Spain
- ³⁰Exercise is Medicine Spain, University of Zaragoza, Zaragoza, Spain
- ³¹The Association for International Sport for All (TAFISA), Frankfurt, Germany
- ³²Sportsmedical Centre Bern-Ittigen, Ittigen, Switzerland
- ³³Sport and Exercise Medicine Switzerland (SEMS), Bern, Switzerland
- ³⁴Shaare Zedek Medical Center, Hebrew University, Jerusalem, Israel
- ³⁵Exercise is Medicine Israel, Hebrew University, Jerusalem, Israel
- ³⁶Centre for Exercise Science and Sports Medicine, University of Witwatersrand, Johannesburg, South Africa
- ³⁷South African Sports Medicine Association (SASMA), Pretoria, South Africa
- ³⁸Unit of Endocrinology - Department of Movement, Human and Health Sciences, University of Rome Foro Italico, Rome, Italy
- ³⁹House of Sport, Europe Active, Brussels, Belgium
- ⁴⁰Department of Sports Medicine, University Grenoble Alpes, Grenoble, Auvergne-Rhône-Alpes, France
- ⁴¹French Society of Exercise and Sports Medicine, Société Française de Médecine de l'Exercice et du Sport, Paris, France
- ⁴²National University of Physical Education and Sport of Ukraine, Kyiv, Ukraine
- ⁴³Department of Sport Medicine and Functional Explorations, University-Hospital (CHU), G. Montpied Hospital, Clermont-Ferrand, France
- ⁴⁴UMR 1019, INRAE, French National Research Institute for Agriculture, Food and Environment, Clermont-Ferrand, France
- ⁴⁵Sports and Exercise Medicine Division, Department of Medicine, Università degli Studi di Padova, Padova, Italy
- ⁴⁶Exercise is Medicine Italy, Università degli Studi di Padova, Padova, Italy
- ⁴⁷German Association of Paediatric and Adolescent Care Specialists, BVKJ - Berufsverband der Kinder- und Jugendärzte, Cologne, Germany
- ⁴⁸Sports Medicine and Pediatrics, Nationwide Children's Hospital and The Ohio State University, Columbus, Ohio, USA
- ⁴⁹American College of Sports Medicine, Indianapolis, Indiana, USA
- ⁵⁰Faculty of Sport and Exercise Sciences, University of Rome 'Foro Italico', Roma, Lazio, Italy
- ⁵¹Hospital de Clínicas José San Martín, University of Buenos Aires, Buenos Aires, Argentina
- ⁵²Australasian College of Sport and Exercise Physicians, Melbourne, Victoria, Australia
- ⁵³AUT Sports Performance Research Institute New Zealand, Auckland, New Zealand
- ⁵⁴German Society of Nephrology (Deutsche Gesellschaft für Nephrologie (DGN)), Berlin, Germany
- ⁵⁵German Society for Neurology (Deutsche Gesellschaft für Neurologie (DGN)), Berlin, Germany
- ⁵⁶Cyprus Olympic Committee, Nicosia, Cyprus
- ⁵⁷Sports Medicine, Swiss Olympic Medical Center, Hopital de la Tour, Meyrin, Geneva, Switzerland
- ⁵⁸SportAdo Consultation - Multidisciplinary Unit of Adolescent Health, University Hospital of Lausanne, Lausanne, Switzerland
- ⁵⁹Department of Health and Human Performance - Facultad de CC de la Actividad Física y del Deporte, INEF Universidad Politécnica de Madrid, Madrid, Spain
- ⁶⁰Senat, Freie und Hansestadt Hamburg, Hamburg, Germany
- ⁶¹European Association of Preventive Cardiology (EAPC), European Society of Cardiology (ECS), Biot, France
- ⁶²Department of Prevention and Sports Medicine, Klinikum rechts der Isar, Technische Universität München, München, Germany
- ⁶³German Diabetes Foundation, Deutsche Diabetes Stiftung, Düsseldorf, Germany
- ⁶⁴Health Research Institute, University of Limerick, Limerick, Ireland
- ⁶⁵Center for Brain and Health Sciences, Aomori University, Aomori, Japan
- ⁶⁶German Olympic Sports Confederation, Deutscher Olympischer Sportbund, Frankfurt am Main, Germany
- ⁶⁷Institute of Sports and Sport Science, University Heidelberg, Heidelberg, Germany
- ⁶⁸Deutscher Verband für Gesundheitssport und Sporttherapie e.V. (DVGS), Hamburg, Germany
- ⁶⁹Sports Medicine, Australian Institute of Sport, Canberra, Canberra, Australia
- ⁷⁰Australian Institute of Sport, Australian Sports Commission, Canberra, Canberra, Australia
- ⁷¹Department of Orthopaedics, University of Illinois at Chicago, Chicago, Illinois, USA
- ⁷²American College of Sports Medicine Foundation, Indianapolis, Indiana, USA
- ⁷³European Federation of Sports Medicine Associations (EFSMA), Lausanne, Switzerland
- ⁷⁴Carol Davila University of Medicine and Pharmacy, Bucharest, Bucharest, Romania
- ⁷⁵Section Sports Medicine, University of Pretoria Faculty of Health Sciences, Pretoria, Gauteng, South Africa
- ⁷⁶Department of Sports Medicine, Medical University of Lodz, Lodz, Poland
- ⁷⁷Moving Medicine, Faculty of Sport and Exercise Medicine UK, Edinburgh, UK
- ⁷⁸Bundesvereinigung Prävention und Gesundheitsförderung e.V. (BVBG), Bonn, Germany
- ⁷⁹Health Science Center, Kansai Medical University, Osaka, Japan
- ⁸⁰Exercise is Medicine Japan, Japanese Society of Physical Fitness and Sports Medicine, Osaka, Japan
- ⁸¹Rīga Stradiņš University, Riga, Latvia
- ⁸²German Society of Orthopaedics and Trauma (Deutsche Gesellschaft für Orthopädie und Unfallchirurgie (DGOU)) with the German Society for Trauma Surgery (DGU) and German Society of Orthopaedics and Orthopaedic Surgery (DGOOC), Berlin, Germany
- ⁸³Platform on Nutrition and Physical Activity, Plattform Ernährung und Bewegung e.V. (peb), Berlin, Germany
- ⁸⁴Exercise is Medicine Norway, Oslo, Norway
- ⁸⁵School of Science and Health, University of Western Sydney, Sydney, New South Wales, Australia
- ⁸⁶Exercise is Medicine Greece, National and Kapodistrian University of Athens, Athens, Greece
- ⁸⁷School of Exercise Science and Dietetics, University of Thessaly, Trikala, Greece
- ⁸⁸German Diabetes Association (Deutsche Diabetes Gesellschaft (DDG)), Berlin, Germany
- ⁸⁹Institute of Epidemiology, Biostatistics and Prevention, Zuerich University, Zuerich, Switzerland
- ⁹⁰German Diabetes Support (diabetesDE - Deutsche Diabetes-Hilfe), Charlottenburg, Germany
- ⁹¹German Alliance for Baths, Bäderallianz Deutschland, Köln, Germany
- ⁹²International Association for Sport and Leisure Facilities, Köln, Germany
- ⁹³German Cardiac Society (Deutsche Gesellschaft für Kardiologie – Herz- und Kreislaufforschung (DGK)), Düsseldorf, Nordrhein-Westfalen, Germany
- ⁹⁴Hamburg Sport Association (Hamburger Sportbund e.V.), Hamburg, Germany
- ⁹⁵UMR INSERM 1272 Hypoxie et poumon, Université Sorbonne Paris Nord - Campus de Bobigny, Bobigny, France
- ⁹⁶Hôpital Jean-Verdier, Médecine de l'exercice et du sport, Assistance Publique - Hôpitaux de Paris, Bondy, France
- ⁹⁷German Society for Physiotherapy Science (Deutsche Gesellschaft für Physiotherapiewissenschaft (DGPTW)), Hamburg, Germany
- ⁹⁸Dept. of Rehabilitation, Saitama Medical University, Saitama, Japan
- ⁹⁹D.A.CH Society Prevention of Cardiovascular Diseases, D.A.CH-Gesellschaft Prävention von Herz-Kreislauf-Erkrankungen, Hamburg, Germany
- ¹⁰⁰University Sports Complex, Institute for Physical Education and Sport, Msida, Malta
- ¹⁰¹World Olympians Association (WOA), Lausanne, Switzerland
- ¹⁰²Global Esports Federation (GEF), Singapore
- ¹⁰³European College of Sport Science, Köln, Germany
- ¹⁰⁴Istanbul Spor Etkinlikleri ve İletmeciliği A S, City of Istanbul, Istanbul, Turkey
- ¹⁰⁵Institute of Sport Sciences and Physiotherapy, Faculty of Medicine, University of Tartu, Tartu, Estonia
- ¹⁰⁶Exercise is Medicine Poland, Legionowo, Poland

- ¹⁰⁷German Hypertension League (Deutsche Hochdruckliga e.V. (DHL)), Heidelberg, Baden-Württemberg, Germany
- ¹⁰⁸German Society for Hypertension and Prevention (Deutsche Gesellschaft für Hypertonie und Prävention), Heidelberg, Germany
- ¹⁰⁹Research Education Centre, ASPETAR - Qatar Orthopaedic and Sports Medicine Hospital, Doha, Qatar
- ¹¹⁰Department of Educational Assistance, Physical and Health Education, Vytautas Magnus University, Vilnius, Lithuania
- ¹¹¹Department of Movement Science, University of Hamburg, Hamburg, Germany
- ¹¹²German Medical Association (Bundesärztekammer), Berlin, Germany
- ¹¹³Sports Laboratory, Sports Medicine and Physical Health Centre, Riga, Latvia, Riga, Latvia
- ¹¹⁴Latvian Sports Medicine Association, Riga, Latvia
- ¹¹⁵Family Medicine, Kaiser Permanente, Fontana, California, USA
- ¹¹⁶Exercise is Medicine Portugal, Universidade de Lisboa, Lisboa, Portugal
- ¹¹⁷Exercise and Health Laboratory, CIPER, Faculdade de Motricidade Humana, Universidade de Lisboa, Lisboa, Portugal
- ¹¹⁸German Society of General Practice and Family Medicine (Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin (DEGAM)), Berlin, Germany
- ¹¹⁹Department of General Practice and Primary Care, University Medical Center, Hamburg, Germany
- ¹²⁰International Society for Physical Activity and Health (ISPAH), Vancouver, British Columbia, Canada
- ¹²¹Society for Orthopaedic and Traumatologic Sports Medicine (GOTS), Jena, Germany
- ¹²²Exercise is Medicine Singapore, Singapore
- ¹²³Department of Sport & Exercise Medicine, Changi General Hospital, Singapore
- ¹²⁴Division of Sports and Exercise Medicine, Department of Sport, Exercise and Health, Basel, Switzerland
- ¹²⁵German Society for the Prevention and Rehabilitation of Cardiovascular Diseases (Deutsche Gesellschaft für Prävention und Rehabilitation von Herz-Kreislaufkrankungen (DGPR)), Koblenz, Germany
- ¹²⁶German Society of Sports Science, Deutsche Vereinigung für Sportwissenschaft (DVS), Frankfurt, Germany
- ¹²⁷Health through Physical Activity, Lifestyle and Sport (HPALS) Research Centre, University of Cape Town, Cape Town, South Africa
- ¹²⁸Federal Institute for Sports Science (Bundesinstitut für Sportwissenschaft (BISp)), Bonn, Nordrhein-Westfalen, Germany
- ¹²⁹Institute for Applied Training Science Leipzig, Leipzig, Sachsen, Germany
- ¹³⁰Division of Surgery and Interventional Science, University College London, London, UK
- ¹³¹Public Health and Family Medicine, University of Western Ontario Schulich School of Medicine and Dentistry, London, Ontario, Canada
- ¹³²Sports Medicine Department, Faculty of Medicine, Ankara University, Ankara, Ankara, Turkey
- ¹³³Department of Sports Medicine and Rehabilitation, Institute of Clinical Medicine, University of Tartu, Tartu, Estonia
- ¹³⁴Department of Medicine, Università degli Studi di Padova, Padova, Italy
- ¹³⁵Laboratory of Regenerative Medicine, Haikou, Hainan, China
- ¹³⁶International Olympic Committee, Lausanne, Switzerland
- ¹³⁷Division of Orthopedic Surgery, University of Oslo, Oslo, Norway
- ¹³⁸Department of Movement, Human and Health Sciences, University of Rome "Foro Italico", Rome, Italy
- ¹³⁹School of Sport and Health Sciences, University of Brighton, Eastbourne, UK

Twitter Boris Gojanovic @DrSportSante, Sebastien Racinais @ephysiol, Jane Thornton @JaneSThornton and Evert Verhagen @evertverhagen

Acknowledgements The authors would like to thank Leonie Fink, Division of Sports and Rehabilitation Medicine, Department of Internal Medicine, Ulm University Hospital, Ulm, Germany.

Contributors The manuscript was drafted by JS, YPP, WB, WvM, FP, TP, CK and JW. The manuscript was then sent out to signatories of the declaration for consent. All coauthors read the manuscript, made recommendations and proved per mail their co-authorship and the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Map disclaimer The inclusion of any map (including the depiction of any boundaries therein), or of any geographic or locational reference, does not imply

the expression of any opinion whatsoever on the part of BMJ concerning the legal status of any country, territory, jurisdiction or area or of its authorities. Any such expression remains solely that of the relevant source and is not endorsed by BMJ. Maps are provided without any warranty of any kind, either express or implied.

Competing interests EV is Editor in Chief of *BMJ Open Sports & Exercise Medicine*.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Jürgen M Steinacker <http://orcid.org/0000-0001-8901-9450>
 Carolin Knoke <http://orcid.org/0000-0002-4907-4428>
 Jose-Antonio Casajus Mallen <http://orcid.org/0000-0002-7215-6931>
 Demetri Constantinou <http://orcid.org/0000-0002-3363-7695>
 Stephane Doutreleau <http://orcid.org/0000-0001-8816-0356>
 Martine Duclos <http://orcid.org/0000-0002-7158-386X>
 Matthew Payton Herring <http://orcid.org/0000-0002-6835-5321>
 Dina Christina Janse van Rensburg <http://orcid.org/0000-0003-1058-6992>
 Susi Kriemler <http://orcid.org/0000-0002-3384-7940>
 François J Lhuissier <http://orcid.org/0000-0003-1491-5896>
 Robert Sallis <http://orcid.org/0000-0001-7633-7345>
 Luis B Sardinha <http://orcid.org/0000-0002-6230-6027>
 Benedict Tan <http://orcid.org/0000-0002-1762-849X>
 Jane Thornton <http://orcid.org/0000-0002-3519-7101>
 Bulent Ulkar <http://orcid.org/0000-0002-4656-8705>
 Evert Verhagen <http://orcid.org/0000-0001-9227-8234>
 Yannis P Pitsiladis <http://orcid.org/0000-0001-6210-2449>

REFERENCES

- López-Valenciano A, Mayo X, Liguori G, *et al*. Changes in sedentary behaviour in European Union adults between 2002 and 2017. *BMJ Public Health* 2020;20:1206.
- World Health Organization. Global status report on physical activity 2022; 2022.
- Wilson F, Gormley J, Hussey J. Exercise therapy in the management of musculoskeletal disorders. In: *Exercise Therapy in the Management of Musculoskeletal Disorders*. London: Wiley-Blackwell, 25 March 2011.
- Pedersen BK, Saltin B. Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015;25 Suppl 3:1-72.
- Tremblay MS, Aubert S, Barnes JD, *et al*. Sedentary behavior research network (SBRN) - terminology consensus project process and outcome. *Int J Behav Nutr Phys Act* 2017;14:75.
- World Health Organization. WHO guidelines on physical activity and sedentary behaviour: web annex: evidence Profiles 2020; 2020.
- Guthold R, Stevens GA, Riley LM, *et al*. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1.9 million participants. *Lancet Glob Health* 2018;6:e1077-86.
- Anderson E, Durstine JL. Physical activity, exercise, and chronic diseases: A brief review. *Sports Med Health Sci* 2019;1:3-10.
- Piepoli MF, Hoes AW, Agewall S, *et al*. European guidelines on cardiovascular disease prevention in clinical practice: the sixth joint task force of the European society of cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of 10 societies and by invited Experts) Developed with the special contribution of the European Association for cardiovascular prevention & rehabilitation (EACPR). *Eur Heart J* 2016;37:2315-81.
- Lancaster GI, Febbraio MA. The Immunomodulating role of exercise in metabolic disease. *Trends Immunol* 2014;35:262-9.
- Liu Y, Yan T, Chu JM-T, *et al*. The beneficial effects of physical exercise in the brain and related pathophysiological mechanisms in neurodegenerative diseases. *Lab Invest* 2019;99:943-57.

- 12 Kramer AF, Hahn S, Cohen NJ, *et al.* Ageing, fitness and Neurocognitive function. *Nature* 1999;400:418–9.
- 13 Buffart LM, Kalker J, Sweegers MG, *et al.* Effects and Moderators of exercise on quality of life and physical function in patients with cancer: an individual patient data meta-analysis of 34 Rcts. *Cancer Treat Rev* 2017;52:91–104.
- 14 Tison GH, Barrios J, Avram R, *et al.* Worldwide physical activity trends since COVID-19 onset. *Lancet Glob Health* 2022;10:e1381–2.
- 15 Mandolesi L, Polverino A, Montuori S, *et al.* Effects of physical exercise on cognitive functioning and wellbeing: biological and psychological benefits. *Front Psychol* 2018;9:509.
- 16 Penedo FJ, Dahn JR. Exercise and well-being: a review of mental and physical health benefits associated with physical activity. *Curr Opin Psychiatry* 2005;18:189–93.
- 17 Naci H, Salcher-Konrad M, Dias S, *et al.* How does exercise treatment compare with antihypertensive medications? A network meta-analysis of 391 randomised controlled trials assessing exercise and medication effects on systolic blood pressure. *Br J Sports Med* 2019;53:859–69.
- 18 Wen CP, Wai JPM, Tsai MK, *et al.* Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* 2011;378:1244–53.
- 19 Onerup A, Arvidsson D, Blomqvist Å, *et al.* Physical activity on prescription in accordance with the Swedish model increases physical activity: a systematic review. *Br J Sports Med* 2019;53:383–8.
- 20 Swedish National Institute of Public Health. Physical activity in the prevention and treatment of disease (FYSS book). In: *Stockholm, Sweden: Professional associations for physical activity, sweden. yrkesföreningar för fysisk aktivitet, yfa*, 2010.
- 21 Carbone S, Del Buono MG, Ozemek C, *et al.* Obesity, risk of diabetes and role of physical activity, exercise training and cardiorespiratory fitness. *Progress in Cardiovascular Diseases* 2019;62:327–33.
- 22 Paffenbarger RS, Wing AL, Hyde RT. Physical activity as an index of heart attack risk in college alumni. *Am J Epidemiol* 1978;108:161–75.
- 23 Sigal RJ, Kenny GP, Boulé NG, *et al.* Effects of aerobic training, resistance training, or both on Glycemic control in type 2 diabetes: a randomized trial. *Ann Intern Med* 2007;147:357–69.
- 24 Ambrosetti M, Abreu A, Corrà U, *et al.* Secondary prevention through comprehensive cardiovascular rehabilitation: from knowledge to implementation. 2020 update. A position paper from the secondary prevention and rehabilitation section of the European Association of preventive cardiology. *Eur J Prev Cardiol* 2021;28:460–95.
- 25 Winzer EB, Woitek F, Linke A. Physical activity in the prevention and treatment of coronary artery disease. *J Am Heart Assoc* 2018;7:e007725.
- 26 Kirsten J, Wais V, Schulz SWW, *et al.* n.d. Sarcopenia screening allows identifying high-risk patients for Allogenic stem cell transplantation. *Cancers*;13:1771.
- 27 Duret C, Breuckmann P, Louchart M, *et al.* Adapted physical activity in community-dwelling adults with neurological disorders: design and outcomes of a fitness-center based program. *Disabil Rehabil* 2022;44:536–41.
- 28 OECD and WHO. Organisation for economic Co-operation and Development (OECD) and world health organization. In: *Step Up! Tackling the Burden of Insufficient Physical Activity in Europe*. Paris: OECD Publishing, 2023.
- 29 Santos AC, Willumsen J, Meheus F, *et al.* The cost of inaction on physical inactivity to public health-care systems: a population-attributable fraction analysis. *Lancet Glob Health* 2023;11:e32–9.
- 30 World Health Organization. *Global action plan on physical activity 2018–2030: more active people for a healthier world*. Geneva: World Health Organization, 2018.
- 31 Sallis R, Young DR, Tartof SY, *et al.* Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48 440 adult patients. *Br J Sports Med* 2021;55:1099–105.
- 32 Medicine Io. Health and Behavior. *The Interplay of Biological, Behavioral, and Societal Influences*. Washington, DC: The National Academies Press, 2001: 395.
- 33 Swinburn B, Vandevijvere S, Kraak V, *et al.* Monitoring and Benchmarking government policies and actions to improve the Healthiness of food environments: a proposed government healthy food environment policy index. *Obes Rev* 2013;14 Suppl 1:24–37.
- 34 Klepac Pogrmilovic B, Ramirez Varela A, Pratt M, *et al.* National physical activity and sedentary behaviour policies in 76 countries: availability, comprehensiveness, implementation, and effectiveness. *Int J Behav Nutr Phys Act* 2020;17:116.
- 35 Deslippe AL, Soanes A, Bouchaud CC, *et al.* Barriers and Facilitators to diet, physical activity and lifestyle behavior intervention adherence: a qualitative systematic review of the literature. *Int J Behav Nutr Phys Act* 2023;20:14.
- 36 Gelius P, Messing S, Forberger S, *et al.* The added value of using the HEPA PAT for physical activity policy monitoring: a four-country comparison. *Health Res Policy Syst* 2021;19:22.
- 37 Gelius P, Messing S, Goodwin L, *et al.* What are effective policies for promoting physical activity? A systematic review of reviews. *Prev Med Rep* 2020;18:101095.
- 38 Lakerveld J, Woods C, Hebestreit A, *et al.* Advancing the evidence base for public policies Impacting on dietary behaviour, physical activity and sedentary behaviour in Europe: the policy evaluation network promoting a Multidisciplinary approach. *Food Policy* 2020;96:101873.
- 39 Dzau VJ, Balatbat CA, Ellaissi WF. Revisiting academic health sciences systems a decade later: discovery to health to population to society. *Lancet* 2021;398:2300–4.
- 40 Smith PJ, Merwin RM. The role of exercise in management of mental health disorders: an integrative review. *Annu Rev Med* 2021;72:45–62.
- 41 Lavie CJ, Johannsen N, Swift D, *et al.* Exercise is medicine - the importance of physical activity, exercise training, cardiorespiratory fitness and obesity in the prevention and treatment of type 2 diabetes. *Eur Endocrinol* 2014;10:18–22.
- 42 Sharpe PA, Burroughs EL, Granner ML, *et al.* Impact of a community-based prevention marketing intervention to promote physical activity among middle-aged women. *Health Educ Behav* 2010;37:403–23.
- 43 Budzynski-Seymour E, Milton K, Mills H, *et al.* A rapid review of communication strategies for physical activity guidelines and physical activity promotion: A review of worldwide strategies. *J Phys Act Health* 2021;18:1014–27.
- 44 Deverka PA, Lavallee DC, Desai PJ, *et al.* Stakeholder participation in comparative effectiveness research: defining a framework for effective engagement. *J Comp Eff Res* 2012;1:181–94.
- 45 Goodman MS, Sanders Thompson VL. The science of Stakeholder engagement in research: classification, implementation, and evaluation. *Transl Behav Med* 2017;7:486–91.
- 46 Concannon TW, Meissner P, Grunbaum JA, *et al.* A new Taxonomy for Stakeholder engagement in patient-centered outcomes research. *J Gen Intern Med* 2012;27:985–91.
- 47 Ghanvatkar S, Kankanhalli A, Rajan V. User models for personalized physical activity interventions: Scoping review. *JMIR Mhealth Uhealth* 2019;7:e11098.
- 48 Breuer E, Lee L, De Silva M, *et al.* Using theory of change to design and evaluate public health interventions: a systematic review. *Implement Sci* 2016;11:63.
- 49 Elwyn G, Frosch D, Thomson R, *et al.* Shared decision making: a model for clinical practice. *J GEN INTERN MED* 2012;27:1361–7.
- 50 Global alliance against physical inactivity. The Hamburg declaration 2021. Available: <https://www.sports-medicine-health-summit.de/en/a-global-alliance-against-physical-inactivity.html> [Accessed 17 Nov 2022].
- 51 International Olympic Committee. IOC and WHO unite further to promote health through sport: key objectives of IOC and WHO memorandum of understanding. 2022. Available: <https://olympics.com/en/news/ioc-world-health-organization-who-unite-health-sport>
- 52 International Olympic Committee. Olympicday Letsmove - move your body and dedicate 30 minutes on June 23rd to get active with Olympians. 2023. Available: <https://olympics.com/en/olympic-day/>
- 53 International Federation of Sports Medicine. FIMS collaborating centres of sports medicine. 2022. Available: <https://www.fims.org/about/ccsm>
- 54 Hollands GJ, Shemilt I, Marteau TM, *et al.* Altering micro-environments to change population health behaviour: towards an evidence base for choice architecture interventions. *BMC Public Health* 2013;13:1218.
- 55 Lobczowska K, Banik A, Romaniuk P, *et al.* Frameworks for implementation of policies promoting healthy nutrition and physically active lifestyle: systematic review. *Int J Behav Nutr Phys Act* 2022;19:16.
- 56 Wong THF, Tapper NJ, Luby SP. Planetary health approaches for dry cities: water quality and heat mitigation. *BMJ* 2020:m4313.
- 57 World Health Organization. *Promoting physical activity through primary health care: a toolkit*. Geneva: World Health Organization, 2021.
- 58 Zukowska J, Gobis A, Krajewski P, *et al.* Which transport policies increase physical activity of the whole of society? A systematic review. *Journal of Transport & Health* 2022;27:101488.
- 59 Keszyüs D, Lauer R, Traub M, *et al.* Effects of statewide health promotion in primary schools on children's sick days, visits to a

- physician and parental absence from work: a cluster-randomized trial. *BMC Public Health* 2016;16:1244.
- 60 Faskunger J. Promoting active living in healthy cities of Europe. *J Urban Health* 2013;90 Suppl 1(Suppl 1):142–53.
- 61 Stout NL, Baima J, Swisher AK, *et al.* A systematic review of exercise systematic reviews in the cancer literature (2005–2017). *PM R* 2017;9:S347–84.
- 62 Fiorito G, Pedron S, Ochoa-Rosales C, *et al.* The role of epigenetic clocks in explaining educational inequalities in mortality: A Multicohort study and meta-analysis. *J Gerontol A Biol Sci Med Sci* 2022;77:1750–9.
- 63 Futreal PA, Coin L, Marshall M, *et al.* A census of human cancer genes. *Nat Rev Cancer* 2004;4:177–83.
- 64 Wahl S, Drong A, Lehne B, *et al.* Epigenome-wide Association study of body mass index, and the adverse outcomes of Adiposity. *Nature* 2017;541:81–6.
- 65 Muniz-Pardos B, Shurlock J, Debruyne A, *et al.* Collateral health issues derived from the COVID-19 pandemic. *Sports Med Open* 2020;6:35.
- 66 The Lancet. COVID-19: a new lens for non-communicable diseases. *Lancet* 2020;396:S0140-6736(20)31856-0.
- 67 Clavel N, Badr J, Gautier L, *et al.* Knowledge and behaviors of general and high-risk adult populations towards COVID-19: A systematic Scoping review. *Public Health Rev* 2021;42:1603979.
- 68 Louro A, Marques da Costa N, Marques da Costa E. Sustainable urban mobility policies as a path to healthy cities—the case study of LMA. *Sustainability* 2019;11:2929.
- 69 Steinacker JM, Matits L, Wendt J, *et al.* Health in all policies? Rethinking prevention. *Dtsch Z Sportmed* 2022;73:3–6.
- 70 McPhail S, Schippers M. An evolving perspective on physical activity counselling by medical professionals. *BMC Fam Pract* 2012;13:31.
- 71 Neunhaeuserer D, Niebauer J, Degano G, *et al.* Sports and exercise medicine in Europe and the advances in the last decade. *Br J Sports Med* 2021;55:1122–4.
- 72 Rose G, Khaw K-T, Marmot MG. Rose's strategy of preventive medicine. In: *Strategy of Preventive Medicine*. Oxford University Press, 10 January 2008.
- 73 Active Well-being Initiative (AWI). Active well-being initiative. 2023. Available: <http://activewellbeing.org/>
- 74 Faculty of Sport and Exercise Medicine. Moving Medicine Birmingham, Available: <https://movingmedicine.ac.uk/> [Accessed 17 Oct 2022].
- 75 Bouma A, van Nassau F, Nauta J, *et al.* Implementing exercise = medicine in routine clinical care; needs for an online tool and key decisions for implementation of exercise = medicine within two Dutch academic hospitals. *BMC Med Inform Decis Mak* 2022;22:250.
- 76 Sallis JF, Cerin E, Kerr J, *et al.* Built environment, physical activity, and obesity: findings from the International physical activity and environment network (IPEN) adult study. *Annu Rev Public Health* 2020;41:119–39.
- 77 Gambhir SS, Ge TJ, Vermesh O, *et al.* Toward achieving precision health. *Sci Transl Med* 2018;10:430.
- 78 Zhang X, Speakman JR. Genetic factors associated with human physical activity: are your genes too tight to prevent you exercising? *Endocrinology* 2019;160:840–52.
- 79 Sharma A, Mentz RJ, Granger BB, *et al.* Utilizing mobile technologies to improve physical activity and medication adherence in patients with heart failure and diabetes mellitus: rationale and design of the TARGET-HF-DM trial. *American Heart Journal* 2019;211:22–33.
- 80 Woessner MN, Tacey A, Levinger-Limor A, *et al.* The evolution of technology and physical inactivity: the good, the bad, and the way forward. *Front Public Health* 2021;9:655491.