



A Century of Change towards Prevention and Minimal Intervention in Cariology



Journal of Dental Research
2019, Vol. 98(6) 611–617
© International & American Associations
for Dental Research 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/0022034519837252
journals.sagepub.com/home/jdr



N.P.T. Innes¹, C.H. Chu², M. Fontana³, E.C.M. Lo², W.M. Thomson⁴,
S. Uribe⁵, M. Heiland⁶, S. Jepsen⁷, and F. Schwendicke⁸

Abstract

Better understanding of dental caries and other oral conditions has guided new strategies to prevent disease and manage its consequences at individual and public health levels. This article discusses advances in prevention and minimal intervention dentistry over the last century by focusing on some milestones within scientific, clinical, and public health arenas, mainly in cariology but also beyond, highlighting current understanding and evidence with future prospects. Dentistry was initially established as a surgical specialty. Dental caries (similar to periodontitis) was considered to be an infectious disease 100 years ago. Its ubiquitous presence and rampant nature—coupled with limited diagnostic tools and therapeutic treatment options—meant that these dental diseases were managed mainly by excising affected tissue. The understanding of the diseases and a change in their prevalence, extent, and severity, with evolutions in operative techniques, technologies, and materials, have enabled a shift from surgical to preventive and minimal intervention dentistry approaches. Future challenges to embrace include continuing the dental profession's move toward a more patient-centered, evidence-based, less invasive management of these diseases, focused on promoting and maintaining oral health in partnership with patients. In parallel, public health needs to continue to, for example, tackle social inequalities in dental health, develop better preventive and management options for existing disease risk groups (e.g., the growing aging population), and the development of reimbursement and health outcome models that facilitate implementation of these evolving strategies. A century ago, almost every treatment involved injections, a drill or scalpel, or a pair of forceps. Today, dentists have more options than ever before available to them. These are supported by evidence, have a minimal intervention focus, and result in better outcomes for patients. The profession's greatest challenge is moving this evidence into practice.

Keywords: caries, minimal intervention dentistry, diagnosis, dental care, evidence based dentistry, public health

Introduction

The concept of minimal (or minimum) intervention dentistry (MID) within oral health care (Kearns et al. 2015) has moved from a fringe topic, taken seriously by only a few, to the center of oral health care. The advances in understanding of dental diseases, human behavior, diagnostics, biomaterials, clinical operative techniques, and technologies have all contributed to our understanding of MID as a patient-centered, biological, and economic paradigm and a contemporary way to deliver dental care. This article focuses on the gradual shift from surgical to minimal intervention and preventive dentistry over the past century and the implications of this shift for public health. We present some MID milestones within the scientific, clinical, and public health arenas and consider the future prospects of MID.

Minimal Intervention Dentistry

For most of human history, the cornerstones of dentistry have been 1) removing carious enamel and dentin (Oxilia et al. 2015), 2) excising infected periodontal tissues, and 3) extracting teeth (and sometimes replacing them). With great foresight, in 1896 G.V. Black expressed a hope that “the day is surely coming, when we will be engaged in practising preventive,

rather than reparative dentistry” (Joseph 2005). Yet, establishing the dental profession as a surgical specialty in the late 19th and early 20th centuries seems to have set the path for a mainly operative approach toward managing dental diseases. This surgical approach was initially grounded in the necessity to treat rampant caries, periodontal disease, and associated pain or infection with very limited means available. The growing

¹School of Dentistry, University of Dundee, Dundee, UK

²Faculty of Dentistry, University of Hong Kong, Hong Kong

³Cariology and Restorative Sciences, School of Dentistry, University of Michigan, Ann Arbor, MI, USA

⁴Department of Oral Sciences, School of Dentistry, University of Otago, Dunedin, New Zealand

⁵School of Dentistry, Universidad Austral de Chile, Valdivia, Chile

⁶Department of Oral and Maxillofacial Surgery, Charité-Universitätsmedizin Berlin, Berlin, Germany

⁷Periodontology, Operative, and Preventive Dentistry, University of Bonn, Bonn, Germany

⁸Operative and Preventive Dentistry, Charité-Universitätsmedizin Berlin, Berlin, Germany

Corresponding Author:

N.P.T. Innes, School of Dentistry, University of Dundee, Park Place, Dundee, DD1 4HR, UK.

Email: n.p.innes@dundee.ac.uk




Figure. Past, present, and future aspects of prevention and minimal intervention in cariology.

understanding of dental caries and, simultaneously, periodontitis as lifestyle-mediated biofilm diseases led to a feeling of futility associated with simply trying to “fix” symptoms without managing the disease or its causes. Improved diagnostics, operative techniques, and biomaterials led to the emergence of novel concepts for preventing and controlling dental caries, followed by a change toward minimally interventive approaches toward other oral and dental conditions (Fig.).

Dental Caries

The recognition of the role of bacteria as a cause of “fermentation” leading to dissolution of tooth substance (Miller 1890) led to the idea that dental caries was an infectious disease, requiring “excision” of affected tissues. Generations of dentists were taught a highly invasive, operatively based approach to managing carious lesions (Black 1908), and this prevailed for almost a century. All contaminated (previously known as “infected”) or demineralized (previously known as “affected”) dental tissue was excised. The appreciation that certain bacteria, notably *Streptococcus mutans*, were more commonly associated with the disease (Keyes 1960; Loesche 1986) supported this approach, as did early successes in managing nondental infectious diseases with antibiotics or vaccination. For dental caries, these approaches (i.e., managing caries as an infectious disease) largely failed to yield significant individual or public health benefits, as evidenced by the widespread experience of rampant caries until the 1980s in most high-income countries.

Over the past 100 years, the futility of this traditionally surgically focused approach has become acknowledged (Fejerskov 2004). Alongside this, a growing recognition of the restorative “spiral” (Elderton 1990) and the escalating invasiveness of retreatments, initiated by placement of the first restoration, is

increasingly seen as part of the problem, rather than the solution for managing caries. In contrast, an emerging understanding of the complexity of dental biofilm (Costerton 1995)—supported by modern analytic technologies, such as genomics, microbiomics, and metabolomics—has facilitated a change in the approach to dental caries management. Knowledge of bacterial species’ interdependence and communication systems (Kolenbrander et al. 2010) and the role of extracellular matrices (Koo et al. 2013) has clarified how bacteria need particular conditions, like the population of a city, to thrive (Marsh 2005). The shift between stages of physiologic biofilm conditions and dysbiosis is a response to environmental pressures (Neilands et al. 2014), a concept that invites management focusing on rebalancing and modulating the biofilm composition (Marsh 2006) and activity and not attempting to eradicate the biofilm per se.

New technologies for detecting and treating dental caries were developed in parallel. These fueled a change from managing the signs of the disease through excision and restoration toward preventing it or controlling its activity. Adhesive dentistry, initiated by the introduction of enamel acid etching and resin bonding to dental tissues (Buonocore 1955), enabled dentists for the first time to remove only the tissue affected by bacterial contamination, instead of cutting cavities according to material demands (e.g., following the cavity preparation rules that G.V. Black had introduced for dental amalgam restorations). The ability to detect lesions at earlier stages with radiography followed later by other technologies may have originally driven earlier intervention to manage the disease at enamel and precavitation stages (Innes and Schwendicke 2017); however, there is a trend that this is reversing, and ultimately, early detection has enabled targeted, less invasive management of early-stage disease.

There was a slow shift toward rebalancing the oral biofilm composition and activity (Massler 1967; Handelman et al. 1976; Elderton 1985; Walsh and Brostek 2013). The emerging health concept of modulating microbiomes via probiotics (i.e., live microorganisms conveying health benefits) has begun to be applied to managing dental caries and periodontal disease oral biofilms (Mira 2018). Probiotics can replace pathogenic bacteria (e.g., *S. mutans*), modulating pathogenicity or altering the resulting immune response. However, their efficacy and the sustainability of any effect for caries prevention and management remains debatable. Notably, most probiotic bacteria are themselves acidogenic and aciduric (e.g., these properties are part of the mechanism that contributes to their health benefits, when applied to the skin or in the gut). This may, in part, explain the heterogeneity in findings from clinical studies of caries prevention and management with probiotics.

Contemporary dentistry has turned towards strategies to arrest or even heal carious lesions. The success of this approach is illustrated by the dramatic decline in caries among children in most high-income countries (Lagerweij and van Loveren 2015). Use of fluoride in general and especially the regular use of fluoride toothpaste for preventing dental caries and arresting carious lesions are supported by strong and consistent evidence (Marinho et al. 2003; Walsh et al. 2010; Wong et al. 2011; Marinho et al. 2013). In fact, combined home-based toothbrushing with fluoride toothpaste is likely to have had the greatest contribution to the reduction in dental caries prevalence and experience (Bratthall et al. 1996).

There have been developments of alternative/supplemental strategies for lesion prevention/mineralization (Ten Cate 2012; Fontana 2016; Featherstone et al. 2018). While calcium-based strategies still have inconsistent evidence for their effectiveness (Slayton et al. 2018; Urquhart et al. 2018), others seem promising, such as fluoride combined with antimicrobials (e.g., stannous fluoride and silver diamine fluoride) or with arginine (Wolff and Schenkel 2018). Novel remineralization methods are showing promise, such as the use of peptides to enhance deeper remineralization (Alkilzy et al. 2018).

Although the direct causal link between sugar and caries development was established through studies as far back as the 1950s and 1960s—the Vipeholm (Gustafsson et al. 1954), Hopewood House (Harris 1963), Tristan da Cunha (Holloway et al. 1962), and Turku sugar studies (Scheinin et al. 1976)—sugar has only recently become of serious interest within dentistry. This omission may have been supported by a range of underlying agendas (Kearns et al. 2015). Only lately has the addictive potential of sugar begun to be understood, with sweet foods' stimulation of the human reward system encouraging repeated overconsumption. In addition, there is only recent acknowledgment of the role of sugar in a range of health conditions, acting through modification of the microbiome and inflammasome. The robust evidence linking the frequency and amount of sugar intake to caries increment and the recognition of the sugar “epidemic” as a problem for broader health, with sugar being a common risk factor for several important non-communicable diseases, have led to prioritization of approaches

to reduce sugar consumption. However, there is still limited evidence to support strategies to promote behavior change at an individual level (Harris et al. 2012; Albino and Tiwari 2016). Hence, public health efforts (e.g., reducing access to sugar-sweetened beverages, reformulating foods and drinks, and sugar taxation) have increasingly become the focus (Schwendicke et al. 2016), and dentistry is increasingly involved in advocating for these measures. Linking our preventive efforts with those of other health disciplines will likely be to the benefit of our patients, the wider public, and our profession (GBD 2015 Risk Factors Collaborators 2016; FDI World Dental Federation 2018; NCD Alliance 2018; World Health Organization 2018).

Restorative materials to seal carious lesions are another group of strategies that modify the environment and thereby the microbiome composition, aiming to arrest the lesion (Handelman et al. 1976). Studies on sealing sound and later carious tissue showed that sealants impede acid diffusion into—and mineral diffusion out of—the dental tissue and also isolate sealed bacteria from their dietary carbohydrate source (Griffin et al. 2008). These studies were initially carried out for enamel and later for noncavitated lesions with the use of sealant materials (Wright et al. 2016; Slayton et al. 2018) and then cavitated ones extending into dentin with the use of more mechanically robust materials (Mertz-Fairhurst et al. 1987). Sealing lesions is less mechanically destructive and more protective of the dental pulp than techniques involving removing all carious tissue (Mertz-Fairhurst et al. 1987; Ricketts et al. 2013). The understanding that bacteria can be sealed (Oong et al. 2008) was the pillar for less invasive carious tissue removal prior to placing a restoration, as cariogenic bacteria and carious tissue were allowed to be left and sealed in proximity to the pulp. Concepts such as stepwise or selective removal of carious tissue are built on this foundation, reducing the risk of pulp exposure and its sequelae. For primary teeth, the Hall technique combines sealing of carious tissue with restoration with a stainless-steel crown (Schwendicke et al. 2016).

Prevention and Minimally Invasive Therapy: Concepts Crossing Disciplines

Over the past century and alongside or following the changes toward minimally invasive treatments for caries (Table), other dental disciplines have adopted the concepts of prevention and MID. These include periodontology and oral and maxillofacial surgery.

The science of periodontology, for example, and the concepts of periodontal etiopathogenesis have evolved from the early descriptions of “alveolar pyorrhea” to the current concepts of “microbial dysbiosis.” During this time, the clinical discipline of periodontics has seen many paradigm changes (Heitz-Mayfield and Lang 2013). Until the 1960s, removal of diseased tissues was considered necessary, and this led to invasive surgical interventions with removal of gingival tissue and/or bone. Later, surgical pocket elimination became the main

Table. Milestones in the Development of Cariology.

Year	Development
1890	Nonspecific plaque hypothesis described by Miller. Periodontal disease described as pyorrhea by Riggs.
1895	Roentgen discovers x-rays. Two weeks later (January 1896), Walkhoff takes the first dental x-ray. Roentgen wins the first Nobel Prize in Physics (1901).
1896	G.V. Black describes classical cavity preparations.
1900	Focal theory: teeth are the reason for systemic infections.
1909	Role of fluoride in fluorosis established.
1925	Raper introduces the bitewing technique.
1945	Grand Rapids initiates water fluoridation for caries prevention.
1948	Introduction of panoramic radiography.
1954	Buonocore introduces acid-etching technique, allowing adhesive dentistry.
1950s and 1960s	Central role of sugar established through studies such as the Vipeholm, Hopewood House, and Tristan da Cunha.
1965	The association between plaque and periodontal disease is established by Loe.
1970s	Costerton develops concepts of the biofilm and complex biofilm communities. Sealing in caries first demonstrated as viable strategy.
1976	Loesche describes the specific plaque hypothesis.
1983	Development of cone beam computed tomography. Hafferjee and Socransky describe clusters of periodontal pathogens.
1980s	Machine-driven periodontal instrumentation introduced.
1987	Intraoral digital radiography introduced.
1990-2000s	Genetic links in periodontitis and caries established.
1991	Marsh describes the ecologic plaque hypothesis.
2018	At the University of Beijing and the Fourth Military Medical University Stomatological Hospital in China, a robot performs a surgical implant with only human programming prior to the procedure.
2018	Machine learning systems allow detection of carious lesions, periodontal bone loss, and apical lesions with accuracies similar or superior to those of experienced dentists.

objective of periodontal therapy, and either gingivectomies or apically positioned flap procedures were widely undertaken. In the 1980s, with more knowledge of periodontal biology, pathogenesis, and wound healing, the necessity for pocket elimination was challenged. The focus shifted to surgical flap procedures enabling access to the root surfaces for scaling and root planing. At the same time, it was demonstrated that non-surgical periodontal therapy was effective even in deep pockets (Badersten et al. 1984), and the concepts of intentional gingival curettage and excessive removal of contaminated cementum were abandoned. A critical probing depth was determined, above which periodontal surgery led to more pocket reduction and clinical attachment gain than scaling and root planning (Lindhe et al. 1982). It was also established that the long-term success of periodontal therapy was critically dependent on the quality of maintenance care and plaque control (Axelsson and Lindhe 1981; Ramfjord et al. 1987). Important advances were made in regenerative periodontal surgery for advanced intrabony defects. Minimally invasive procedures show advantages in wound-healing outcomes, recession, and patient morbidity (Cortellini and Tonetti 2009; Trombelli et al. 2012). Twenty-year outcomes of regenerative therapy are promising (Cortellini et al. 2017). Novel periodontal tissue bioengineering is also under development (Fretwurst et al. 2018).

Similar to caries, the concepts of primary and secondary prevention are crucial in periodontology (Tonetti et al. 2015). Periodontitis is preventable through effective management of gingivitis and promotion of healthy lifestyles at both the population and individual levels (Chapple et al. 2015; Jepsen et al. 2017). Risk profiling and stratification are of key importance (Giannobile et al. 2013). Overwhelming evidence shows that,

for the majority of patients, periodontitis can be treated and effectively managed by a series of sequential phases of care, with appropriate, sustained changes to self-care and smoking habits (Tonetti et al. 2017a).

Oral and maxillofacial surgery has also become less invasive, with comparable reductions in morbidity. Instead of open approaches, techniques with endoscopes, microscopes, or robotic systems are now routine in many procedures, such as those involving the sinuses, temporomandibular joint, or salivary glands, but also in tumor surgery and some aesthetic procedures. As an example, the field of transoral robotic surgery has seen dynamic development during the last years, with the number of articles published rising from 3 in 2006 to 123 in 2016 alone (Poon et al. 2018). Minimal invasive surgery can also involve virtual planning based on 3-dimensional image data and their transfer via individual drill guides, robotic systems, patient-specific implants, or navigation systems (Heiland et al. 2004). However, these systems need further refinement. Notably, though, these strategies require more preoperative data, possibly involving greater radiation exposure, and are more expensive than conventional approaches. More robust evidence of the benefits to patients is also needed.

Implications and the Future

In most high-income countries, there have been improvements in dental health. Dental caries experience among children has been declining for decades (although early childhood caries remains a problem), and a simultaneous reduction has been seen recently among adults and seniors (Lagerweij and van Loveren 2015). More teeth are being retained by adults and

seniors than ever before, with the need for removable prostheses decreasing dramatically. Despite the retention of more teeth, the number of periodontally affected teeth does not necessarily seem to be increasing, possibly due to falling smoking rates (Haisman-Welsh and Thomson 2012). While it would be pretentious to assume that these successes are grounded only in the changing approach of the profession toward dental health—from surgery to MID and prevention—this shift in managing dental disease has certainly contributed to it. Arising from these successes, however, are implications for how we manage dental caries and other conditions. Future shifts in morbidity and population demographics will further affect the direction of our profession. A number of aspects need highlighting, and these are considered in turn.

First, dentistry is becoming more complex. Fifty years ago, a dentist would usually be presented with a fairly homogeneous group of patients: most were seen every 6 months, had cavitated lesions, required restorations (mainly amalgam), and lost teeth early on. With the differential and age- and socially specific decline in the number of carious, restored, or missing teeth, dentists are nowadays faced with a highly heterogeneous clientele. Hence, there is a need for more targeted diagnostics and personalized management to ensure that each of these very different patients receives the best treatment. So far, the tools for identifying the specific preventive and therapeutic needs of a patient rely largely on history taking and clinical and/or radiographic findings. Most caries risk assessment systems, for example, use a range of risk indicators (caries experience, dietary habits, oral hygiene, fluoride intake), weight them, and then assign the individual's risk status. The same is true for periodontal risk assessment tools and the recent reclassification of periodontal diseases, with multidimensional staging and grading (Papapanou et al. 2018). Based on such risk assessment or classification, active and supportive care can be determined with treatment thresholds (Schwendicke 2018). However, most risk assessment or disease classification systems have been only sparsely validated, show limited accuracy, and are not truly “personalized” but allow only a rough stratification of individuals according to risk. It can be assumed that—with progress in systems medicine allowing deeper insights into individual disease mechanisms based on clinical, imagery, sample (saliva or blood), or routine (also nonhealth) data—new insights into individual risk and current health conditions will be possible. Digital technologies will enable the best use of these data, eventually paving the way for “4P dentistry”: precision, personalized, preventive, and participatory dental care (Hood and Flores 2012). Such an approach promises considerable health gains at an individual level. However, at a population level, it could increase inequality, as those with the disease are less likely to seek or afford care (Knight and Thomson 2018).

This leads to the second aspect: while dentistry has changed for the better, not everyone has benefited unequivocally from improvements in oral health. Its nature as chronic, cumulative behavior-mediated diseases mean that dentistry shares the problem of inequality with other noncommunicable diseases, many of which disproportionately affect those of lower

social position. This is associated with numerous social, structural, and/or institutional factors. There is compelling evidence that dental services utilization, for example, is also highly unfairly distributed among different social, ethnic, economic, and educational groups (Reda et al. 2018). Dental care is therefore unlikely to reduce inequalities in health but may even accentuate them. Increasing the application of public health policies—such as promoting fluoridation, antismoking policies, or healthy diets—may help to reduce such social inequality. Moreover, preventive care and MID need to be made available to billions of people currently without access to conventional dental care, such as through the atraumatic restorative technique (Frencken et al. 2012) or interprofessional collaborations.

Third, with older people living longer and retaining more teeth but having multiple morbidities, the focus of dental care will need to change. Cumulative (lifelong) caries experience is unlikely to reduce (Broadbent et al. 2013; Knight and Thomson 2018) but be shifted into older age (a phenomenon known as “morbidity compression”). Periodontal diseases will be mostly affected, and this need will have to be addressed among elderly individuals (Schwendicke et al. 2018; Tonetti et al. 2017b). Prevention and MID will need to account for this, with new concepts and products being required specifically for older people.

Finally, disease detection and assessment need to adapt. The concept that lesions can be active or inactive means that the “scar” of the disease (e.g., a discolored inactive lesion or a residual radiographic radiolucency) is not necessarily a problem or a health risk for the patient. This is an important conceptual hurdle for the profession because the traditional approach has been to aim for a fully restored dentition as the gold standard. However, contemporary dentistry often involves neither removal of carious lesions (because this would do more harm than good, as discussed earlier) nor healing of carious lesions (because we are currently unable to do so); thus, it is vital to be able to categorize a lesion as “arrested” or “active” and to monitor this. Only a few validated systems to assess lesion activity are available; these include the International Caries Detection and Assessment System (Pitts 2004) and the Nyvad criteria (Nyvad and Baelum 2018). Moreover, the radiographic detection of carious lesions has been found to be limited in accuracy. Proximal radiographic examination has low sensitivity to detect small lesions (Schwendicke et al. 2015) and low agreement when used by general dentists. Machine learning—in particular, the application of deep convolutional neural networks to build predictive models for radiographic imagery data—may help to improve accuracy in diagnosis.

Conclusion

In dentistry and specifically cariology, we take for granted many of the things that we now do on a daily basis, and it is easy to forget that there was little understanding of the pathogenesis of dental caries and other conditions 100 years ago, with almost every treatment involving injections, a drill, a scalpel, or a pair of forceps. Today, we have far better insight into

the etiopathology of dental caries as a biofilm-based but behavior-mediated disease. Dentists have a larger number of treatment options and evidence available to them than ever before, and, in parallel with other oral conditions, are moving toward more minimal-intervention, evidence-supported, but personalized treatment options that focus on promoting and maintaining oral health. Preventive and public health efforts across the globe have achieved a great deal in this, although the benefits are not yet universal.

We nevertheless face many challenges, some of which we have discussed here. However, one of the greatest challenges lies not in future developments, as important as these are, but in implementing what we know is most effective and making the best use of what we have already developed and available to us: translating the science, ideas, and concepts of cariology and minimally invasive dentistry into practice.

Author Contributions

N.P.T. Innes, F. Schwendicke, S. Jepsen, M. Heiland, S. Uribe, contributed to conception and design, drafted and critically revised the manuscript; C.H. Chu, M. Fontana, E.C.M. Lo, W.M. Thomson, contributed to conception and design, critically revised the manuscript. All authors gave final approval and agree to be accountable for all aspects of the work.

Acknowledgments

The authors' institutions supported this work. There were no external sources of funding. The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article.

ORCID iDs

N.P.T. Innes  <https://orcid.org/0000-0002-9984-0012>
S. Uribe  <https://orcid.org/0000-0003-0684-2025>

References

- Albino J, Tiwari T. 2016. Preventing childhood caries: a review of recent behavioral research. *J Dent Res*. 95(1):35–42.
- Alkilzy M, Santamaria RM, Schmoedel J, Splieth CH. 2018. Treatment of carious lesions using self-assembling peptides. *Adv Dent Res*. 29(1):42–47.
- Axelsson P, Lindhe J. 1981. Effect of controlled oral hygiene procedures on caries and periodontal disease in adults: results after 6 years. *J Clin Periodontol*. 8(3):239–248.
- Badersten A, Nilveus R, Egelberg J. 1984. Effect of nonsurgical periodontal therapy: II. Severely advanced periodontitis. *J Clin Periodontol*. 11(1):63–76.
- Black GV. 1908. The technical procedures in filling teeth: a work on operative dentistry. Chicago (IL): Medico-Dental Publishing.
- Brathall D, Hansel-Petersson G, Sundberg H. 1996. Reasons for the caries decline: what do the experts believe? *Eur J Oral Sci*. 104(4, Pt 2):416–422.
- Broadbent JM, Foster Page LA, Thomson WM, Poulton R. 2013. Permanent dentition caries through the first half of life. *Br Dent J*. 215(7):E12.
- Buonocore MG. 1955. A simple method of increasing the adhesion of acrylic filling materials to enamel surfaces. *J Dent Res*. 34(6):849–853.
- Chapple IL, Van der Weijden F, Doerfer C, Herrera D, Shapira L, Polak D, Madianos P, Louropoulou A, Machtei E, Donos N, et al. 2015. Primary prevention of periodontitis: managing gingivitis. *J Clin Periodontol*. 42 Suppl 16:S71–S76.
- Cortellini P, Buti J, Pini Prato G, Tonetti MS. 2017. Periodontal regeneration compared with access flap surgery in human intra-bony defects 20-year follow-up of a randomized clinical trial: tooth retention, periodontitis recurrence and costs. *J Clin Periodontol*. 44(1):58–66.
- Cortellini P, Tonetti MS. 2009. Improved wound stability with a modified minimally invasive surgical technique in the regenerative treatment of isolated interdental intrabony defects. *J Clin Periodontol*. 36(2):157–163.
- Costerton JW. 1995. Overview of microbial biofilms. *J Ind Microbiol*. 15(3):137–140.
- Elderton RJ. 1985. Management of early dental caries in fissures with fissure sealant. *Br Dent J*. 158(7):254–258.
- Elderton RJ. 1990. Clinical studies concerning re-restoration of teeth. *Adv Dent Res*. 4:4–9.
- FDI World Dental Federation. 2018. Oral health and noncommunicable diseases; [accessed 2018 Oct 12]. <https://www.fdiworlddental.org/what-we-do/advocacy/oral-health-and-noncommunicable-diseases>.
- Featherstone JD, Fontana M, Wolff M. 2018. Novel anticaries and remineralization agents: future research needs. *J Dent Res*. 97(2):125–127.
- Fejerskov O. 2004. Changing paradigms in concepts on dental caries: consequences for oral health care. *Caries Res*. 38(3):182–191.
- Fontana M. 2016. Enhancing fluoride: clinical human studies of alternatives or boosters for caries management. *Caries Res*. 50 Suppl 1:22–37.
- Frencken JE, Leal SC, Navarro MF. 2012. Twenty-five-year atraumatic restorative treatment (ART) approach: a comprehensive overview. *Clin Oral Invest*. 16(5):1337–1346.
- Fretwurst T, Larsson L, Yu SH, Pilipchuk SP, Kaigler D, Giannobile WV. 2018. Periodontal tissue bioengineering: is the future now? *Compend Contin Educ Dent*. 39(4):218–223.
- GBD 2015 Risk Factors Collaborators. 2016. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the global burden of disease study 2015. *Lancet*. 388(10053):1659–1724.
- Giannobile WV, Braun TM, Caplis AK, Doucette-Stamm L, Duff GW, Korman KS. 2013. Patient stratification for preventive care in dentistry. *J Dent Res*. 92(8):694–701.
- Griffin SO, Oong E, Kohn W, Vidakovic B, Gooch BF; CDC Dental Sealant Systematic Review Work Group, Bader J, Clarkson J, Fontana MR, Meyer DM, Rozier RG, et al. 2008. The effectiveness of sealants in managing caries lesions. *J Dent Res*. 87(2):169–174.
- Gustafsson BE, Quensel CE, Lanke LS, Lundqvist C, Grahnen H, Bonow BE, Krasse B. 1954. The vipeholm dental caries study; the effect of different levels of carbohydrate intake on caries activity in 436 individuals observed for five years. *Acta Odontol Scand*. 11(3–4):232–264.
- Haisman-Welsh RJ, Thomson WM. 2012. Changes in periodontitis prevalence over two decades in New Zealand: evidence from the 1988 and 2009 national surveys. *N Z Dent J*. 108(4):134–138.
- Handelman SL, Washburn F, Wopperer P. 1976. Two-year report of sealant effect on bacteria in dental caries. *J Am Dent Assoc*. 93(5):967–970.
- Harris R. 1963. Biology of the children of Hopewood House, Bowral, Australia: 4. Observations on dental-caries experience extending over five years (1957–61). *J Dent Res*. 42:1387–1399.
- Harris R, Gamboa A, Dailey Y, Ashcroft A. 2012. One-to-one dietary interventions undertaken in a dental setting to change dietary behaviour. *Cochrane Database Syst Rev*. (3):CD006540.
- Heiland M, Habermann CR, Schmelzle R. 2004. Indications and limitations of intraoperative navigation in maxillofacial surgery. *J Oral Maxillofac Surg*. 62(9):1059–1063.
- Heitz-Mayfield LJ, Lang NP. 2013. Surgical and nonsurgical periodontal therapy: learned and unlearned concepts. *Periodontol*. 62(1):218–231.
- Holloway PJ, James PM, Slack GL. 1962. Dental caries among the inhabitants of Tristan da Cunha: (d). The last count. *R Soc Health J*. 82:139.
- Hood L, Flores M. 2012. A personal view on systems medicine and the emergence of proactive p4 medicine: predictive, preventive, personalized and participatory. *N Biotechnol*. 29(6):613–624.
- Innes NPT, Schwendicke F. 2017. Restorative thresholds for carious lesions: systematic review and meta-analysis. *J Dent Res*. 96(5):501–508.
- Jepsen S, Blanco J, Buchalla W, Carvalho JC, Dietrich T, Dorfer C, Eaton KA, Figuero E, Frencken JE, Graziani F, et al. 2017. Prevention and control of dental caries and periodontal diseases at individual and population level: consensus report of group 3 of joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *J Clin Periodontol*. 44 Suppl 18:S85–S93.
- Joseph R. 2005. The father of modern dentistry—Dr. Greene Vardiman Black (1836–1915). *J Conserv Dent*. 8(2):5–6.
- Kearns CE, Glantz SA, Schmidt LA. 2015. Sugar industry influence on the scientific agenda of the National Institute of Dental Research's 1971 National Caries Program: a historical analysis of internal documents. *PLoS Med*. 12(3):e1001798.

- Keyes PH. 1960. The infectious and transmissible nature of experimental dental caries: findings and implications. *Arch Oral Biol.* 1:304–320.
- Knight ET, Thomson WM. 2018. A public health perspective on personalized periodontics. *Periodontol* 2000. 78(1):195–200.
- Kolenbrander PE, Palmer RJ Jr, Periasamy S, Jakubovics NS. 2010. Oral multi-species biofilm development and the key role of cell-cell distance. *Nat Rev Microbiol.* 8(7):471–480.
- Koo H, Falsetta ML, Klein MI. 2013. The exopolysaccharide matrix: a virulence determinant of cariogenic biofilm. *J Dent Res.* 92(12):1065–1073.
- Lagerweij MD, van Loveren C. 2015. Declining caries trends: are we satisfied? *Curr Oral Health Rep.* 2(4):212–217.
- Lindhe J, Socransky SS, Nyman S, Haffajee A, Westfelt E. 1982. “Critical probing depths” in periodontal therapy. *J Clin Periodontol.* 9(4):323–336.
- Loesche WJ. 1986. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev.* 50(4):353–380.
- Marinho VC, Higgins JP, Sheiham A, Logan S. 2003. Fluoride toothpastes for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev.* (1):CD002278.
- Marinho VC, Worthington HV, Walsh T, Clarkson JE. 2013. Fluoride varnishes for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev.* (7):CD002279.
- Marsh PD. 2005. Dental plaque: biological significance of a biofilm and community life-style. *J Clin Periodontol.* 32 Suppl 6:7–15.
- Marsh PD. 2006. Dental plaque as a biofilm and a microbial community—implications for health and disease. *BMC Oral Health.* 6 Suppl 1:S14.
- Massler M. 1967. Pulpal reactions to dental caries. *Int Dent J.* 17(2):441–460.
- Mertz-Fairhurst EJ, Call-Smith KM, Shuster GS, Williams JE, Davis QB, Smith CD, Bell RA, Sherrer JD, Myers DR, Morse PK, et al. 1987. Clinical performance of sealed composite restorations placed over caries compared with sealed and unsealed amalgam restorations. *J Am Dent Assoc.* 115(5):689–694.
- Miller WD. 1890. The micro-organisms of the human mouth. Philadelphia (PA): The S. S. White Dental Mfg. Co.
- Mira A. 2018. Oral microbiome studies: potential diagnostic and therapeutic implications. *Adv Dent Res.* 29(1):71–77.
- NCD Alliance. 2018. FDI World Dental Federation; [accessed 2018 Oct 12]. <https://ncdalliance.org/who-we-are/ncd-alliance-structure/supporters-consultation-group/fdi-world-dental-federation>.
- Neilands J, Troedsson U, Sjodin T, Davies JR. 2014. The effect of delmopinol and fluoride on acid adaptation and acid production in dental plaque biofilms. *Arch Oral Biol.* 59(3):318–323.
- Nyvad B, Baelum V. 2018. Nyvad criteria for caries lesion activity and severity assessment: a validated approach for clinical management and research. *Caries Res.* 52(5):397–405.
- Oong EM, Griffin SO, Kohn WG, Gooch BF, Caufield PW. 2008. The effect of dental sealants on bacteria levels in caries lesions: a review of the evidence. *J Am Dent Assoc.* 139(3):271–278.
- Oxilia G, Peresani M, Romandini M, Matteucci C, Spiteri CD, Henry AG, Schulz D, Archer W, Crezzini J, Boschin F, et al. 2015. Earliest evidence of dental caries manipulation in the late upper palaeolithic. *Sci Rep.* 5:12150.
- Papapanou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, Flemmig TF, Garcia R, Giannobile WV, Graziani F, et al. 2018. Periodontitis: consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-implant Diseases and Conditions. *J Periodontol.* 89 Suppl 1:S173–S182.
- Pitts N. 2004. “ICDAS”—an international system for caries detection and assessment being developed to facilitate caries epidemiology, research and appropriate clinical management. *Community Dent Health.* 21(3):193–198.
- Poon H, Li C, Gao W, Ren H, Lim CM. 2018. Evolution of robotic systems for transoral head and neck surgery. *Oral Oncol.* 87:82–88.
- Ramfjord SP, Caffesse RG, Morrison EC, Hill RW, Kerry GJ, Appleberry EA, Nissle RR, Stults DL. 1987. 4 modalities of periodontal treatment compared over 5 years. *J Clin Periodontol.* 14(8):445–452.
- Reda SF, Reda SM, Thomson WM, Schwendicke F. 2018. Inequality in utilization of dental services: a systematic review and meta-analysis. *Am J Public Health.* 108(2):e1–e7. doi:10.2105/AJPH.2017.304180.
- Ricketts D, Lamont T, Innes NP, Kidd E, Clarkson JE. 2013. Operative caries management in adults and children. *Cochrane Database Syst Rev.* (3):CD003808.
- Scheinin A, Makinen KK, Ylitalo K. 1976. Turku sugar studies: V. Final report on the effect of sucrose, fructose and xylitol diets on the caries incidence in man. *Acta Odontol Scand.* 34(4):179–216.
- Schwendicke F. 2018. Tailored dentistry: from “one size fits all” to precision dental medicine? *Oper Dent.* 43(5):451–459.
- Schwendicke F, Frencken JE, Bjørndal L, Maltz M, Manton DJ, Ricketts D, Van Landuyt K, Banerjee A, Campus G, Doméjean S, et al. 2016. Managing carious lesions: consensus recommendations on carious tissue removal. *Adv Dent Res.* 28(2):58–67.
- Schwendicke F, Krois J, Kocher T, Hoffmann T, Micheelis W, Jordan RA. 2018. More teeth in more elderly: periodontal treatment needs in Germany 1997–2030. *J Clin Periodontol.* 45(12):1400–1407.
- Schwendicke F, Paris S, Stolpe M. 2015. Detection and treatment of proximal caries lesions: milieu-specific cost-effectiveness analysis. *J Dent.* 43(6):647–655.
- Slayton RL, Urquhart O, Araujo MWB, Fontana M, Guzman-Armstrong S, Nascimento MM, Novy BB, Tinanoff N, Weyant RJ, Wolff MS, et al. 2018. Evidence-based clinical practice guideline on nonrestorative treatments for carious lesions: a report from the American dental association. *J Am Dent Assoc.* 149(10):837–849.e819.
- Ten Cate JM. 2012. Novel anticaries and remineralizing agents: prospects for the future. *J Dent Res.* 91(9):813–815.
- Tonetti MS, Jepsen S, Jin L, Otomo-Corgel J. 2017a. Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: A call for global action. *J Clin Periodontol.* 44(5):456–462.
- Tonetti MS, Bottenberg P, Conrads G, Eickholz P, Heasman P, Huysmans MC, Lopez R, Madianos P, Muller F, Needleman I, et al. 2017b. Dental caries and periodontal diseases in the ageing population: call to action to protect and enhance oral health and well-being as an essential component of healthy ageing—consensus report of group 4 of the joint EFP/ORCA workshop on the boundaries between caries and periodontal diseases. *J Clin Periodontol.* 44 Suppl 18:S135–S144.
- Tonetti MS, Chapple IL, Jepsen S, Sanz M. 2015. Primary and secondary prevention of periodontal and peri-implant diseases: introduction to, and objectives of the 11th European Workshop on Periodontology consensus conference. *J Clin Periodontol.* 42 Suppl 16:S1–S4.
- Trombelli L, Simonelli A, Schincaglia GP, Cucchi A, Farina R. 2012. Single-flap approach for surgical debridement of deep intraosseous defects: a randomized controlled trial. *J Periodontol.* 83(1):27–35.
- Urquhart O, Tampi MP, Pilcher L, Slayton RL, Araujo MWB, Fontana M, Guzman-Armstrong S, Nascimento MM, Novy BB, Tinanoff N, et al. 2018. Nonrestorative treatments for caries: systematic review and network meta-analysis. *J Dent Res.* 98(1):14–26.
- Walsh LJ, Brostek AM. 2013. Minimum intervention dentistry principles and objectives. *Aust Dent J.* 58:3–16.
- Walsh T, Worthington HV, Glenny AM, Appelbe P, Marinho VC, Shi X. 2010. Fluoride toothpastes of different concentrations for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev.* (1):CD007868.
- Wolff MS, Schenkel AB. 2018. The anticaries efficacy of a 1.5% arginine and fluoride toothpaste. *Adv Dent Res.* 29(1):93–97.
- Wong MC, Clarkson J, Glenny AM, Lo EC, Marinho VC, Tsang BW, Walsh T, Worthington HV. 2011. Cochrane reviews on the benefits/risks of fluoride toothpastes. *J Dent Res.* 90(5):573–579.
- World Health Organization. 2018. E-library of Evidence for Nutrition Actions (ELENA); [accessed 2018 Oct 12]. <http://www.who.int/elena/titles/free-sugars-adults-ncds/en/>.
- Wright JT, Tampi MP, Graham L, Estrich C, Crall JJ, Fontana M, Gillette EJ, Nový BB, Dhar V, Donly K, et al. 2016. Sealants for preventing and arresting pit-and-fissure occlusal caries in primary and permanent molars: a systematic review of randomized controlled trials—a report of the American Dental Association and the American Academy of Pediatric Dentistry. *J Am Dent Assoc.* 147(8):631–645.e618. Erratum in: *J Am Dent Assoc.* 2017;148(4):210.