

Register-based information on thyroid diseases in Europe: lessons and results from the EUthyroid collaboration

Short title: Thyroid diseases in Europe

Authors

Line Tang Møllehave¹

Marie Holm Eliassen¹

Ieva Strēle²

Allan Linneberg^{1 3}

Rodrigo Moreno-Reyes^{4 5}

Ludmila B. Ivanova⁶

Zvonko Kusić^{7 8}

Iris Erlund⁹

Till Ittermann¹⁰

Ingibjorg Gunnarsdottir^{11 12}

Jonathan Eli Arbelle^{13 14}

Aaron Milton Troen¹⁵

Valdis Pīrāgs¹⁶

Lisbeth Dahl¹⁷

Alicja Hubalewska-Dydejczyk¹⁸

Malgorzata Trofimiuk-Müldner¹⁸

João Jacome de Castro¹⁹

Mafalda Marcelino¹⁹

Simona Gaberšček^{20 21}

Katja Zaltel^{20 21}

¹ Center for Clinical Research and Prevention, Bispebjerg and Frederiksberg Hospital, Frederiksberg, Denmark

² The Institute of Occupational Safety and Environmental Health, Riga Stradiņš University, Riga, Latvia

³ Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

⁴ Nuclear Medicine Department, Erasme Hospital, Brussels, Belgium

⁵ Université Libre de Bruxelles, Belgium

⁶ Faculty of Medicine, Sofia University St. Kl. Ohridski, Sofia, Bulgaria

⁷ Croatian Academy of Sciences and Arts, Zagreb, Croatia

⁸ School of Medicine, Zagreb, Croatia

⁹ Dept. of Government Services, Finnish Institute for Health and Welfare, Helsinki, Finland

¹⁰ Institute for Community Medicine, University Medicine Greifswald, Greifswald, Germany

¹¹ Faculty of Food Science and Nutrition, University of Iceland, Reykjavik, Iceland

¹² Department of Clinical Nutrition, Landspítali-National University Hospital, Reykjavik, Iceland

¹³ Division of Medicine, Maccabi Healthcare Services, Tel Aviv, Israel

¹⁴ Goldman School of Medicine, Ben Gurion University of the Negev, Beer Sheva, Israel

¹⁵ The Institute of Biochemistry Food Science and Nutrition, The Hebrew University of Jerusalem, Rehovot, Israel

¹⁶ Internal Medicine, University of Latvia, Riga, Latvia

¹⁷ Seafood, Nutrition and Environmental State, Institute of Marine Research (IMR), Bergen, Norway

¹⁸ Chair and Department of Endocrinology, Jagiellonian University, Medical College, Cracow, Poland

¹⁹ Endocrine Department, Armed Forces Hospital, Lisbon, Portugal

²⁰ Department of Nuclear Medicine, University Medical Centre Ljubljana, Ljubljana, Slovenia

²¹ Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia

Manuel Puig-Domingo^{22 23}
Sofia Manousou^{24 25}
Helena Filipsson Nyström^{26 27 28}
Michael Bruce Zimmermann²⁹
Karen R. Mullan³⁰
Jayne Valerie Woodside³¹
Henry Völzke¹⁰
Betina Heinsbæk Thuesen¹

Correspondence

Line Tang Møllehave, Center for Clinical Research and Prevention, Bispebjerg and Frederiksberg Hospital, Capital Region, Frederiksberg Hospital, Hovedvejen, Entrance 5, Nordre Fasanvej 57, DK-2000 Frederiksberg, Denmark. Tlf: +45 38633275. E-mail: line.tang.moellehave@regionh.dk.

Keywords: Europe, registries, iodine, thyroid diseases

Words in abstract: 242

Words in manuscript: 3,410

Figures: 1

Tables: 5

References: 52

²² Dpt of Endocrinology and Nutrition, Germans Trias Research Institute and Hospital, Badalona, Spain

²³ Universitat Autònoma de Barcelona, Spain

²⁴ Sahlgrenska Academy, Institute of Medicine, Gothenburg, Sweden

²⁵ Frölunda Specialist Hospital, Västra Frölunda, Sweden

²⁶ Department of Endocrinology, Specialist Medicine, Sahlgrenska University Hospital, Gothenburg, Sweden

²⁷ Department of internal Medicine, Institute of Medicine, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

²⁸ Wallenberg Centre of Molecular and Translational Medicine, Sahlgrenska University Hospital, Gothenburg, Sweden

²⁹ Health Sciences and Technology, ETH Zurich, Zurich, Switzerland

³⁰ Regional Centre for Endocrinology and Diabetes, Belfast Health and Social Care Trust, Belfast, United Kingdom

³¹ Centre for Public Health, Queen's University Belfast, Belfast, United Kingdom

ABSTRACT

Objective: Registers of diagnoses and treatments exist in different forms in the European countries and are potential sources to answer important research questions. Prevalence and incidence of thyroid diseases are highly dependent on iodine intake and, thus, iodine deficiency disease prevention programs. We aimed to collect European register data on thyroid outcomes to compare the rates between countries/regions with different iodine status and prevention programs.

Design: Register-based cross-sectional study.

Methods: National register data on thyroid diagnoses and treatments were requested from 23 European countries/regions. The provided data were critically assessed for suitability for comparison between countries/regions. Sex- and age-standardized rates were calculated.

Results: Register data on ≥ 1 thyroid diagnoses or treatments were available from 22 countries/regions. After critical assessment, data on medication, surgery, and cancer were found suitable for comparison between 9, 10, and 13 countries/regions, respectively. Higher rates of antithyroid medication and thyroid surgery for benign disease and lower rates of thyroid hormone therapy were found for countries with iodine insufficiency before approx. 2001, and no relationship was observed with recent iodine intake or prevention programs.

Conclusions: Collation of register data on thyroid outcomes from European countries is impeded by a high degree of heterogeneity in the availability and quality of data between countries. Nevertheless, a relationship between historic iodine intake and rates of treatments for hyper- and hypothyroid disorders is indicated. This study illustrates both the challenges and the potential for application of register data of thyroid outcomes across Europe.

INTRODUCTION

Register-based data on treatments and diagnoses of thyroid disease can be used as proxies for thyroid disease frequency in a population. This has e.g. been successfully utilized in Denmark (2). In many European countries, treatments for thyroid disease and/or patients diagnosed with thyroid disorders are documented in nationwide or regional registers. This encompasses registers of diseases, medications and other treatments, hospital admissions and ambulatory care, as well as registries related to health insurances. Ideally, for use in research projects, a register should cover the entire country/region or a large, well-defined sub-population whereby it is representative, and the data should be of acceptable validity (3).

European registers have previously been collated for research projects in e.g. melanoma (5), asthma (6), and cerebral palsy (7) and several initiatives to promote and improve the collaboration and utilization of registers in Europe are established (8, 9). However, the availability and applicability of register data for assessment of thyroid outcomes from the European countries have not yet been examined. After thorough assessment of the quality and comparability of data across countries, European registers can potentially represent valuable and efficient data sources for evaluating differences in thyroid outcomes across the countries/regions.

Iodine intake in a population is a main factor behind thyroid disease epidemiology. Iodine deficiency is associated with adverse outcomes ranging from severe mental retardation and stunting to goiter/thyroid nodules and hyperthyroidism (1). However, the window of adequate iodine intake is narrow, hence, iodine excess may increase the risk of hypothyroidism and autoimmune thyroid disorders in a population (10).

Most regions of Europe will be iodine deficient without programs to prevent iodine deficiency disorders (IDD). However, the implementation and type of IDD prevention programs depends on stakeholder barriers, political prioritization (13), and cultural differences. The IDD prevention programs range from general recommendations for intake of iodine-rich foods, over recommendations for iodine supplementation especially for pregnant and lactating women, to voluntary or mandatory iodine fortification (IF), most commonly addition of iodine to table salt and salt in bread (14). An overview of IF programs and iodine intake across 23 European countries/regions is presented in **Table 1**. The optimal level of iodine intake and the most effective IDD prevention program remains to be established.

In many countries, the assessment of median urinary iodine concentration in representative population samples are used to classify the iodine status and to steer IDD prevention programs. While this approach is relatively simple, it is not sufficient to evaluate the aim of such prevention programs, which is to reduce prevalence and incidence of thyroid diseases and related treatments and to save expenditures in health-care systems (4). This can potentially be achieved by analyses of register data. We hypothesized that efficient IDD prevention programs ensuring sufficient iodine intake are robustly associated with a low burden of thyroid related diseases in populations.

The EUthyroid project was an ambitious collaborative project with participation and representatives from most EU countries funded by the European Union's Horizon 2020 research and innovation program under grant agreement no. 634453. The overall aim of the EUthyroid project was to "*evaluate iodine deficiency prevention and monitoring programs in European countries, to initiate capacity building for harmonized European ID prevention and monitoring programs.*"

In this paper we report lessons and results from work package 1 of the EUthyroid project. First, we evaluate the availability, applicability, and comparability of register data on thyroid outcomes in the European

countries, and second, we report country-specific rates of thyroid outcomes and seek to identify patterns in the relationship with iodine supply across Europe.

MATERIALS and METHODS

Register data

Representatives from each country participating in the EUthyroid project were requested to deliver national register data on diagnoses, medical treatments, and procedures for thyroid disorders. All data were requested for the calendar years 2011-2014 and stratified by sex and age in 5-year intervals. If this was not available, we allowed for the available data that best approximated these criteria, e.g. data from 2015, data from a well-defined sub-population, or data in 10-year age intervals. Specific diagnoses and treatments requested were:

Diagnoses: Annual number of persons diagnosed with thyroid cancer (ICD-10: C73), congenital iodine-deficiency syndrome (ICD-10:E00), iodine-deficiency-related thyroid disorders and allied conditions (ICD-10: E01), subclinical iodine-deficiency (ICD-10: E02), other hypothyroidism (ICD-10: E03), other nontoxic goiter (ICD-10: E04), thyrotoxicosis/hyperthyroidism (ICD-10: E05), thyroiditis (ICD-10: E06), other thyroid disorders (ICD-10: E07), and one or more thyroid related diagnoses (ICD-10: E00-E07).

Treatments: Annual number of persons with at least one dispensed dose and total number of defined daily doses (DDD) of antithyroid medication and thyroid hormone therapy. Annual number of persons with at least one procedure and the total number procedures for all thyroid surgeries, thyroid surgeries with benign indication, all radioiodine treatments, radioiodine treatments with benign indication, and fine needle biopsies.

The demographic data of the total populations (total number of individuals and stratified by sex and 5-year age intervals) were derived from Eurostat (<http://ec.europa.eu/eurostat>) or provided from the national EUthyroid representatives for regional or sub-national populations.

Ethical and/or data handling approvals were obtained in each country according to national regulations to obtain aggregated register data. Only aggregated data was collected and, thus, no patient level data (personal data) was handled thereby protecting the privacy the individuals.

Critical assessment and selection of data and variables

The national representatives were also requested to provide a description of the data sources and health care organization factors relevant to assess the completeness and validity of the data provided.

The applicability of the provided data from each country was critically assessed. For inclusion in the database and calculation of raw rates the provided data had to fulfill the criteria for the requested data. For comparison between countries and calculation of standardized rates, the data and variables had to be stratified by sex and age, collected from comparable data sources and registered by comparable methods. The critical assessment was conducted by the senior author, BHT.

Iodine intake

Populations' iodine intakes were derived from published reports of urinary iodine concentrations in school aged children or adults and summarized in **Table 1**. Iodine intake was classified as insufficient, adequate or

excess according to current WHO epidemiological recommendations (15). Early and recent iodine intake were classified as iodine intake approximately 15-20 years before and after 2001, respectively.

Statistical methods

Rates of medications were calculated as one-year period prevalence per 10,000 persons. Rates of diagnoses were calculated per year per 10,000 persons. Raw rates were calculated with the treatments and diagnoses as the numerator and the demographic data of the population as the denominator in the provided sex and/or age strata where possible.

Standardized rates were calculated to remove the effect of the differences in sex and age distributions between the European countries. This was done by calculating the proportion of each sex and age-specific strata in the total European Standard Population (16) and multiplying this by the raw rate in the corresponding strata for each country before summarizing the overall standardized rate in each country. When data were provided for more than one calendar year, the rates were calculated as the average of the calendar years available.

Comparisons between countries were done using standardized rates. No formal statistical tests were applied in the comparison of rates between countries.

RESULTS

Data availability and data assessment

Table 2 provides an overview of the data sources and **Figure 1** provides an overview of the flow of the data acquisition, critical assessments, and calculations.

Register data for at least one thyroid outcome was available and provided from 22 out of 23 participating countries/regions. However, critical assessment concluded that data from 3 countries were considered unsuitable to be included in the database. See overview of data and assessments in the **Supplementary Table S1 and S2**. Raw rates for data included in the database are presented in the **Supplementary Table S3-S5**.

The outcome variables were grouped into 6 categories: thyroid medication, thyroid surgery, radioiodine treatment, fine needle biopsies, diagnoses of thyroid diseases, and diagnoses of incident thyroid cancer. Sex- and age stratified data from all categories were available from 4 countries, whereas data from only one outcome variable was available from 4 countries. The critical assessment of the data included in the database found that data on use of thyroid medication, number of thyroid surgeries and incidence of thyroid cancer were of acceptable quality and thus appropriate for calculation of standardized rates and comparison between countries.

In contrast, data on diagnoses of thyroid diseases except for thyroid cancer were not considered suitable for comparisons across countries. In some countries, registrations included patients treated in primary care whereas only patients attending hospitals were included in other countries, and it was not possible to distinguish between incidence and prevalence data. Furthermore, in some countries registration of diagnoses were based on the ICD-9 system whereas the ICD-10 system was used in other countries and the conversion of data from ICD-9 to ICD-10 caused some misclassification. Data on radioiodine and fine needle biopsies were only available from few countries and were generally considered unreliable.

Thyroid medication

Data from 9 countries were found to be appropriate for comparison of standardized rates of thyroid medication between countries. Hereby we found that the use of antithyroid medication was lowest for Sweden, Finland, Norway and Spain (8.6-13.7 users/10,000 persons) and highest for Denmark, Latvia and Hungary (34.8-43.2 users/10,000 persons). Conversely, the use of thyroid hormone therapy was lowest for Latvia, Denmark and Hungary (192.4-253.1 users/10,000 persons) and highest for Czech Republic, Belgium, Finland, Sweden and Norway (424.8-612.8 users/10,000 persons) (**Table 3**).

There was a clear indication of higher use of antithyroid medication relative to thyroid hormone therapy for countries with early iodine insufficiency. However, there was no apparent pattern in the association between recent iodine intake and use of thyroid medications.

Thyroid surgeries

For thyroid surgeries, data from 10 countries were found to be appropriate for comparison between countries. We found that the standardized rates of thyroid surgeries for benign indications were lowest for Czech Republic, Slovenia, Norway, Sweden and Spain (1.4-2.0 patients/10,000 persons), all with early iodine adequacy, and highest for Belgium, Latvia, and Hungary (4.3-5.1 patients/10,000 persons), all with early iodine insufficiency (**Table 4**). However, no clear pattern was indicated for thyroid surgeries for all indications or when compared with recent iodine intake.

Thyroid cancer

Data on standardized rates of thyroid cancer diagnoses were assessed to be appropriate for comparison between 13 countries/regions. The lowest rates were found in Bulgaria, Northern Ireland, and Scotland (0.4-0.5 patients/10,000 persons) and the highest rates in Denmark, Latvia and Hungary (0.9-1.4 patients/10,000 persons) (**Table 5**), with no clear pattern in the association between early or recent iodine intake and cancer rates.

DISCUSSION

In this first collation of register data on thyroid outcomes from European countries, we found a high degree of heterogeneity in the availability and quality of register data between countries. Data for thyroid medication, thyroid surgery and thyroid cancer were suitable for comparison between 9, 10, and 13 countries, respectively. These data indicated higher rates of antithyroid medication and thyroid surgery for benign disease and lower rates of thyroid hormone therapy in countries with early iodine insufficiency. There was no apparent pattern in the association between thyroid cancer and iodine intake.

Registers, data availability, and data quality

All but one of the 23 participating countries had data on at least one thyroid outcome. However, the data sources, number of variables and quality of register data varied widely due to differences in the organization of the healthcare systems, digitization, history/culture, and political decisions. This is also recognized in previous papers acknowledging the vast potential of the European registers but also addressing the challenges (8, 9, 17).

Nationwide health registries were established in the Scandinavian countries, Latvia, Hungary and in the Region of Catalonia in Spain. In contrast, Croatia, Germany, Greece, Great Britain, North Macedonia, and

Switzerland provided no data from registers fulfilling the requested criteria. Register data from Czech Republic, Belgium, and Bulgaria were derived from health insurance companies covering the majority of the population or a well-defined sub-population. These encompass both public and private health insurance companies. Information on thyroid cancer was available from most countries.

For all data sources the representativity and validity were assessed. There was little concern regarding the representativity in the Nordic registers (18), but some health insurance registers may carry a risk of social stratification hampering the representativity. European cancer registries have a long history and the coverage is generally high (8). The validity of registers could be compromised by over-, under-, or misreporting. Thyroid medications require prescriptions in most European countries and are rarely prescribed in hospitals. This increases the probability of correct reporting as most registers are based on redeemed prescriptions (17). However, e.g. in the Latvian National Health Service register, only reimbursed prescriptions were registered. Some patients may have private insurance or choose to pay fully out-of-pocket and are thus not registered. The validity of thyroid surgeries was also assessed to be appropriate for the included countries. In cancer registries, the data is generally reliable but comparability is still somewhat influenced by diverse methods between countries (8). In contrast, diagnoses other than cancer were considered to be unreliable. Registration of thyroid diagnoses vary both between countries and can also vary within countries between health care providers and specialists. E.g. the rates of congenital iodine-deficiency syndrome were far higher in some countries than what national experts regards as plausible. Validation studies are rarely available for any of the registers (17).

In addition, between-country differences in other risk factors for thyroid outcomes (e.g. alcohol (19, 20), smoking (21), and other micronutrient deficiencies (22)) and between-country differences in diagnostic activity and treatment practices may also influence the rates of thyroid outcomes independently of iodine intake. Especially differences in ultrasound utilization can impact the detection and thus diagnoses and treatment for thyroid nodules and cancer. Despite clinical practice guidelines developed by recognized professional associations, surveys among clinicians find some variability and divergence from guidelines in clinical management of Grave's disease, hypothyroidism, and thyroid nodules (23-25). Furthermore, during 2001-2009/2015 a fall in threshold for treatment initiation of thyroid hormone therapy was observed in both Great Britain and in Denmark leading to more prescriptions of thyroid hormone therapy, but the fall in treatment threshold was markedly steeper in Denmark (26, 27). It is a limitation that we were unable to adjust for these between-country differences.

Overall, collection and comparisons of register-based data from European countries is time consuming and complicated. When it comes to the burden of thyroid diseases, potential consequences of differences in iodine intake status and IDD prevention programs may be overshadowed by within- and between-country sources of bias and confounding.

Differences in thyroid medication, surgery, and cancer rates

Despite the potential biases in comparisons, we still found fairly consistent rates of thyroid outcomes in line with previous research both when comparing the Scandinavian countries who have rather similar registers, healthcare systems, and health behaviors, but also when including data from quite different countries.

Thus, the findings of the highest rates of treatments most frequently applied for hyperthyroidism and/or structural thyroid diseases in countries with early iodine insufficiency and, conversely, the highest rates of treatment for hypothyroidism in countries with early iodine adequacy correspond with previous studies within countries comparing areas with high and low iodine intake (28-30) and over time, when iodine

adequacy was achieved after introduction of IDD prevention programs (31-34). However, it is a limitation that the specific indication for treatment is not known.

We also found that rates of thyroid cancer did not correlate with early or recent iodine intake, for which previous results are not fully consistent (35). Previous studies have indicated that increased iodine intake is associated with an increase in the ratio between papillary and follicular thyroid cancer (36), which was not measured in our study. Nevertheless, whether the overall incidence of thyroid cancer is influenced by the population level iodine intake is still not clear (35).

We chose to only examine overall patterns in the relationship between iodine intake and rates of thyroid outcomes because the available data on iodine intake were collected by different methods, timepoints, and populations across the European countries and thus not sufficiently comparable. If valid data on early and recent iodine intake in Europe are harmonized it will be possible to perform more advanced statistical analyses of associations between the burden of thyroid diseases and iodine intake (37, 38).

Of note, our results indicate a higher correlation between early iodine intake and thyroid outcomes than recent iodine intake. Many of the included countries have introduced IDD prevention programs and achieved iodine sufficiency during the past two decades, and thus the adult population may still be affected by the exposure to decades of iodine insufficiency. Biologically it is plausible that thyroid function and thereby thyroid disease is influenced by the lifetime iodine intake level (39). Furthermore, the introduction of IF can cause a temporary rise in incident hyperthyroidism in countries with previous ID (1) or may induce thyroid dysfunction due to excessive iodine intake (40). This could affect the rates of treatment in several included countries. In Denmark, steady states were achieved for rates of thyrotoxicosis but not for hypothyroidism approximately 10 years after implementation of mandatory IF (33, 34). Our study indicates that it may take even longer before the difference between former iodine adequacy and iodine insufficiency is negated between countries. Despite overall iodine adequacy, iodine deficiency can still be present in vulnerable groups, e.g. pregnant women, which is observed in several countries (37).

The results show no differences between countries characterized by different IDD prevention strategies. Thus, a best-practice model of IDD prevention cannot be identified through this study. This may be explained, firstly, by the difference in coverage and impact of especially voluntary IF. Secondly, IDD prevention programs are optimally designed to compensate for the level of iodine deficiency in the target population, thus, similar iodine intakes may be achieved from different IDD prevention strategies.

Strengths and limitations

The main strengths of this study are, firstly, the strong collaboration between representatives from thyroid research and clinical environments from the European countries. The national EUthyroid representatives had or obtained knowledge of the availability and documentation of data that would not be possible for one researcher from one country. Secondly, the acquisition of several thyroid outcomes enabled us to evaluate different aspects of thyroid disease and the relationship with iodine intake. Thirdly, the critical assessment of data secured that only data of appropriate quality was included. However, the critical assessment was based on evaluations of the available documentation, and thorough validation of most of the variables was not possible. Raw rates were calculated for all provided data included in the database, however, it is important to note that these should be interpreted with caution because the validity of some data is uncertain and comparison of the raw rates will be confounded by differences in sex- and age distributions. Only few variables met the criteria for comparison between countries and these variables were only available for few countries, which is a major limitation. This in combination with the

heterogeneity of iodine intake data and lack of confounding factors limited the generalizability and our ability to draw more firm conclusions on the effect of iodine intake and IDD prevention programs.

This study elucidates the limitations of the registers and points out the need for greater participation and more uniformity in reporting in the future to optimally utilize the valuable registry information. The current increase in digitization and awareness of the possibilities within register research may lead to more valid register data sources on thyroid diagnoses and treatment with easy access and utilization in the European countries in the future. This will be of great value to thyroid research.

Conclusion and perspectives

The availability and quality of register data on thyroid diagnoses and treatments are highly heterogeneous between European countries. Thus, comparison of rates of thyroid outcomes is currently only possible between a limited number of countries. The available data indicate lower rates of antithyroid medication and thyroid surgery for benign disease and higher rates of thyroid hormone therapy in countries with early iodine intake adequacy, and no difference for thyroid cancer or according to recent iodine intake.

As stated in the Krakow Declaration, evaluation of primary thyroid outcomes is essential in the control of IDD prevention programs (41). Registers may be an effective tool for this evaluation. Thus, the present project serves as initial exploration of the availability and quality of register data of thyroid outcomes across Europe, while more in-depth examination of the validity and comparability of registers is still needed to fully utilize the potential of the registers.

DECLARATION OF INTEREST

The authors have no conflicts of interest to declare. The author Alicja Hubalewska-Dydejczyk is an editor of the journal.

FUNDING

This work was supported by the European Union's Horizon 2020 research and innovation program under grant agreement no. 634453. Data analyses were further supported by the project SUPERTHYREOSE, funded by the German G-BA (VSF2_2019-167). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

(42-47) (47-52)

REFERENCES

1. Zimmermann MB, Boelaert K. Iodine deficiency and thyroid disorders. *The lancet Diabetes & endocrinology*. 2015;3(4):286-95.
2. Mollehave LT, Linneberg A, Skaaby T, Knudsen N, Jorgensen T, Thuesen BH. Trends in treatments of thyroid disease following iodine fortification in Denmark: a nationwide register-based study. *Clinical epidemiology*. 2018;10:763-70.
3. Erlangsen A, Fedyszyn I. Danish nationwide registers for public health and health-related research. *Scandinavian journal of public health*. 2015;43(4):333-9.
4. Völzke H, Caron P, Dahl L, de Castro JJ, Erlund I, Gaberšček S, Gunnarsdottir I, Hubalewska-Dydejczyk A, Ittermann T, Ivanova L, et al. Ensuring Effective Prevention of Iodine Deficiency Disorders. *Thyroid : official journal of the American Thyroid Association*. 2016;26(2):189-96.
5. Forsea AM. Melanoma Epidemiology and Early Detection in Europe: Diversity and Disparities. *Dermatology practical & conceptual*. 2020;10(3):e2020033.
6. van Bragt JJMH, Hansen S, Djukanovic R, Bel EHD, ten Brinke A, Wagers SS, Maitland-van der Zee AH, Porsbjerg C. SHARP: Enabling generation of real-world evidence on a pan-European scale to improve the lives of individuals with severe asthma. *ERJ Open Research*. 2021:00064-2021.
7. Alriksson-Schmidt AI, Jeglinsky-Kankainen IFD, Jahnsen RB, Hollung SJ, Andersen GL, Hågglund GV. Flaunting our assets. Making the most of the Nordic registry goldmine: Cerebral palsy as an example. *Scandinavian journal of public health*. 2020;48(1):113-8.
8. Siesling S, Louwman WJ, Kwast A, van den Hurk C, O'Callaghan M, Rosso S, Zanetti R, Storm H, Comber H, Steliarova-Foucher E, et al. Uses of cancer registries for public health and clinical research in Europe: Results of the European Network of Cancer Registries survey among 161 population-based cancer registries during 2010-2012. *European journal of cancer (Oxford, England : 1990)*. 2015;51(9):1039-49.
9. Lovestone S. The European medical information framework: A novel ecosystem for sharing healthcare data across Europe. *Learning health systems*. 2020;4(2):e10214.
10. Laurberg P, Cerqueira C, Ovesen L, Rasmussen LB, Perrild H, Andersen S, Pedersen IB, Carle A. Iodine intake as a determinant of thyroid disorders in populations. *Best practice & research Clinical endocrinology & metabolism*. 2010;24(1):13-27.
11. Rasmussen LB, Ovesen L, Bulow I, Jorgensen T, Knudsen N, Laurberg P, Pertild H. Dietary iodine intake and urinary iodine excretion in a Danish population: effect of geography, supplements and food choice. *The British journal of nutrition*. 2002;87(1):61-9.
12. van der Reijden OL, Zimmermann MB, Galetti V. Iodine in dairy milk: Sources, concentrations and importance to human health. *Best practice & research Clinical endocrinology & metabolism*. 2017;31(4):385-95.
13. Schaffner M, Rochau U, Stojkov I, Qerimi Rushaj V, Völzke H, Marckmann G, Lazarus JH, Oberaigner W, Siebert U. Barriers Against Prevention Programs for Iodine Deficiency Disorders in Europe: A Delphi Study. *Thyroid : official journal of the American Thyroid Association*. 2020.
14. Fortification of food-grade salt with iodine for the prevention and control of iodine deficiency disorders. *World Health Organization* 2014.
15. Zimmermann MB, Andersson M. Assessment of iodine nutrition in populations: past, present, and future. *Nutrition reviews*. 2012;70(10):553-70.
16. Revision of the European Standard Population - Report of Eurostat's task force. Luxembourg: Publications Office of the European Union: European Commission, 2013.
17. Pacurariu A, Plueschke K, McGettigan P, Morales DR, Slattery J, Vogl D, Goedecke T, Kurz X, Cave A. Electronic healthcare databases in Europe: descriptive analysis of characteristics and potential for use in medicines regulation. *BMJ open*. 2018;8(9):e023090.
18. Smith Jervelund S, De Montgomery CJ. Nordic registry data: value, validity and future. *Scandinavian journal of public health*. 2020;48(1):1-4.

19. Carle A, Bulow Pedersen I, Knudsen N, Perrild H, Ovesen L, Rasmussen LB, Jorgensen T, Laurberg P. Graves' hyperthyroidism and moderate alcohol consumption: evidence for disease prevention. *Clinical endocrinology*. 2013;79(1):111-9.
20. Carlé A, Pedersen IB, Knudsen N, Perrild H, Ovesen L, Rasmussen LB, Jørgensen T, Laurberg P. Moderate alcohol consumption may protect against overt autoimmune hypothyroidism: a population-based case-control study. *European journal of endocrinology / European Federation of Endocrine Societies*. 2012;167(4):483-90.
21. Gruppen EG, Kootstra-Ros J, Kobold AM, Connelly MA, Touw D, Bos JHJ, Hak E, Links TP, Bakker SJL, Dullaart RPF. Cigarette smoking is associated with higher thyroid hormone and lower TSH levels: the PREVEND study. *Endocrine*. 2020;67(3):613-22.
22. O'Kane SM, Mulhern MS, Pourshahidi LK, Strain JJ, Yeates AJ. Micronutrients, iodine status and concentrations of thyroid hormones: a systematic review. *Nutrition reviews*. 2018;76(6):418-31.
23. Burch HB, Burman KD, Cooper DS, Hennessey JV, Vietor NO. A 2015 Survey of Clinical Practice Patterns in the Management of Thyroid Nodules. *The Journal of clinical endocrinology and metabolism*. 2016;101(7):2853-62.
24. Burch HB, Burman KD, Cooper DS, Hennessey JV. A 2013 survey of clinical practice patterns in the management of primary hypothyroidism. *The Journal of clinical endocrinology and metabolism*. 2014;99(6):2077-85.
25. Burch HB, Burman KD, Cooper DS. A 2011 survey of clinical practice patterns in the management of Graves' disease. *The Journal of clinical endocrinology and metabolism*. 2012;97(12):4549-58.
26. Taylor PN, Iqbal A, Minassian C, Sayers A, Draman MS, Greenwood R, Hamilton W, Okosieme O, Panicker V, Thomas SL, et al. Falling threshold for treatment of borderline elevated thyrotropin levels-balancing benefits and risks: evidence from a large community-based study. *JAMA internal medicine*. 2014;174(1):32-9.
27. Medici BB, Nygaard B, La Cour JL, Grand MK, Siersma V, Nicolaisdottir DR, Lind B, Olivarius NF, Andersen CL. Changes in prescription routines for treating hypothyroidism between 2001 and 2015 - an observational study of 929,684 primary care patients in Copenhagen. *Thyroid : official journal of the American Thyroid Association*. 2019.
28. Du Y, Gao Y, Meng F, Liu S, Fan Z, Wu J, Sun D. Iodine deficiency and excess coexist in china and induce thyroid dysfunction and disease: a cross-sectional study. *PLoS one*. 2014;9(11):e111937.
29. Pedersen IB, Knudsen N, Jorgensen T, Perrild H, Ovesen L, Laurberg P. Large differences in incidences of overt hyper- and hypothyroidism associated with a small difference in iodine intake: a prospective comparative register-based population survey. *The Journal of clinical endocrinology and metabolism*. 2002;87(10):4462-9.
30. Szabolcs I, Podoba J, Feldkamp J, Dohan O, Farkas I, Sajgó M, Takáts KI, Góth M, Kovács L, Kressinszky K, et al. Comparative screening for thyroid disorders in old age in areas of iodine deficiency, long-term iodine prophylaxis and abundant iodine intake. *Clinical endocrinology*. 1997;47(1):87-92.
31. Baltisberger BL, Minder CE, Burgi H. Decrease of incidence of toxic nodular goitre in a region of Switzerland after full correction of mild iodine deficiency. *European journal of endocrinology / European Federation of Endocrine Societies*. 1995;132(5):546-9.
32. Mostbeck A, Galvan G, Bauer P, Eber O, Atefie K, Dam K, Feichtinger H, Fritzsche H, Haydl H, Köhn H, et al. The incidence of hyperthyroidism in Austria from 1987 to 1995 before and after an increase in salt iodization in 1990. *European journal of nuclear medicine*. 1998;25(4):367-74.
33. Petersen M, Knudsen N, Carle A, Andersen S, Jorgensen T, Perrild H, Ovesen L, Rasmussen LB, Thuesen BH, Pedersen IB. Increased Incidence Rate of Hypothyroidism After Iodine Fortification in Denmark: A 20-Year Prospective Population-Based Study. *The Journal of clinical endocrinology and metabolism*. 2019;104(5):1833-40.

34. Petersen M, Knudsen N, Carle A, Andersen S, Jorgensen T, Perrild H, Ovesen L, Rasmussen LB, Thuesen BH, Pedersen IB. Thyrotoxicosis after iodine fortification. A 21-year Danish population-based study. *Clinical endocrinology*. 2018;89(3):360-6.
35. Zimmermann MB, Galetti V. Iodine intake as a risk factor for thyroid cancer: a comprehensive review of animal and human studies. *Thyroid research*. 2015;8:8.
36. Feldt-Rasmussen U. Iodine and cancer. *Thyroid : official journal of the American Thyroid Association*. 2001;11(5):483-6.
37. Bath SC. The challenges of harmonising the iodine supply across Europe. *The lancet Diabetes & endocrinology*. 2017;5(6):411-2.
38. Ittermann T, Johner S, Below H, Leiterer M, Thamm M, Remer T, Völzke H. Interlaboratory variability of urinary iodine measurements. *Clinical chemistry and laboratory medicine*. 2018;56(3):441-7.
39. Chung HR. Iodine and thyroid function. *Annals of pediatric endocrinology & metabolism*. 2014;19(1):8-12.
40. Farebrother J, Zimmermann MB, Andersson M. Excess iodine intake: sources, assessment, and effects on thyroid function. *Annals of the New York Academy of Sciences*. 2019;1446(1):44-65.
41. The Krakow Declaration on Iodine: Tasks and Responsibilities for Prevention Programs Targeting Iodine Deficiency Disorders. *European thyroid journal*. 2018;7(4):201-4.
42. Nystrom HF, Brantsaeter AL, Erlund I, Gunnarsdottir I, Hulthen L, Laurberg P, Mattisson I, Rasmussen LB, Virtanen S, Meltzer HM. Iodine status in the Nordic countries - past and present. *Food & nutrition research*. 2016;60:31969.
43. Elvbakken KT, Meltzer HM. Research, knowledge, and policy on goitre and iodine in Norway (1850–2016). *Centaurus*. 2021;63(2):396-415.
44. Vandevijvere S, Mourri AB, Amsalkhir S, Avni F, Van Oyen H, Moreno-Reyes R. Fortification of bread with iodized salt corrected iodine deficiency in school-aged children, but not in their mothers: a national cross-sectional survey in Belgium. *Thyroid : official journal of the American Thyroid Association*. 2012;22(10):1046-53.
45. Regulation No. 696: Regulations Regarding Edible Salt, (2015).
46. Global Fortification Data Exchange [updated 17th Feb 2021 accessed 20th Feb 2021]. Available from: <https://fortificationdata.org/map-nutrient-levels-in-fortification-standards/>.
47. Trofimiuk-Müldner M, Konopka J, Sokołowski G, Dubiel A, Kieć-Klimczak M, Kluczyński Ł, Motyka M, Rzepka E, Walczyk J, Sokołowska M, et al. Current iodine nutrition status in Poland (2017): is the Polish model of obligatory iodine prophylaxis able to eliminate iodine deficiency in the population? *Public health nutrition*. 2020;23(14):2467-77.
48. Iodine status worldwide : WHO Global Database on Iodine Deficiency. Geneva: World Health Organization, 2004.
49. Global Scorecard of Iodine Nutrition (2014-2015) and (2020) Iodine Global Network [accessed 20 Feb 2021]. Available from: <https://www.ign.org/scorecard.htm>.
50. Vejbjerg P, Knudsen N, Perrild H, Carle A, Laurberg P, Pedersen IB, Rasmussen LB, Ovesen L, Jorgensen T. Effect of a mandatory iodization program on thyroid gland volume based on individuals' age, gender, and preceding severity of dietary iodine deficiency: a prospective, population-based study. *The Journal of clinical endocrinology and metabolism*. 2007;92(4):1397-401.
51. Mullan K, Hamill L, Doolan K, Young I, Smyth P, Flynn A, Walton J, Meharg AA, Carey M, McKernan C, et al. Iodine status of teenage girls on the island of Ireland. *European Journal of Nutrition*. 2020;59(5):1859-67.
52. Ivanova LB. The Prevention and Control of Iodine Deficiency – Achievements and Problems in Bulgaria. *Endocrinologia*. 2018(3):121-31.

Figure 1: Overview of data availability, critical assessments, and calculation of rates




Figure 1: Overview of data availability, critical assessments, and calculations of rates

338x190mm (96 x 96 DPI)

Table 1: IDD prevention and iodine intake in the European countries


Country	Country code	Iodine fortification strategy (42-47)	Iodine intake, Median UIC, µg/L (population, year) (47-52)
Belgium	BE	Voluntary fortification of salt, 10-15 ppm since 2009	Insufficient, 80 (SAC, 1998) Adequate, 113 (SAC, 2010-2011)
Bulgaria	BG	Mandatory fortification of salt, 28-55 ppm since 1994/1996	Adequate, 111 (SAC, 1996) Adequate, 182 (SAC, 2008) Adequate, 176 (SAC, 2012)
Croatia	HR	Mandatory fortification of salt, 19 ppm since 2011	Adequate, 140 (SAC, 2002) Adequate, 248 (SAC, 2009)
Czech Republic	CZ	Voluntary fortification of salt, 27 ppm since 2016	Adequate, 119 (SAC, 2000) Adequate 163 (SAC, 2010)
Denmark	DK	Mandatory fortification of salt, 13 ppm since 2000	Insufficient, 61 (A, 1997-1998) Adequate, 101 (A, 2004-2005) Adequate, 145 (SAC, 2015)
Finland	FI	Voluntary fortification of salt, 25 ppm since approx. 1946	Adequate, 164 (A, 1997) Insufficient, 83 (A, 2013)
Germany	DE	Voluntary fortification of salt, 20 ppm since 1993	Adequate, 148 (SAC, 1999) Adequate, 122 (SAC, 2003-2006) Insufficient, 89 (SAC+A, 2014-2017)
United Kingdom	UK		
<i>Northern Ireland</i>		None	Adequate, 111 (SAC, 2014-2015)
<i>Scotland</i>		None	N/A
Greece	GR	Voluntary fortification of salt, 30 ppm since 1987	Adequate, 202 (2001) Adequate, 132 (A, 2017-2018)
Hungary	HU	Mandatory fortification of salt, 19 ppm since 2013	Insufficient, 80 (SAC, 1994-1997) Adequate, 228 (SAC, 2005)
Iceland	IS	None	Adequate, 150 (A, 1998) Adequate, 200 (A, 2007-2008)
Ireland	IE	None	Insufficient, 82 (A, 1999) Adequate, 111 (SAC, 2014-2015)
Israel	IL	None	Insufficient, 83 (SAC, 2016)
Latvia	LV	Voluntary fortification of salt, 20-50 (40-100) ppm since 2005	Insufficient, 59 (SAC, 2000) Adequate, 110 (SAC, 2010-2011)
North Macedonia	MK	Mandatory fortification of salt, 25 ppm since 1999	Adequate, 241 (2007) Adequate, 216 (2016)
Norway	NO	Voluntary fortification of salt, 5 ppm since 1938	Adequate, 104 (1999-2001) Insufficient, 75 (A, 2017-2018)
Poland	PL	Mandatory fortification of salt, 23 ppm since 1997	Insufficient, 84 (SAC, 1999) Adequate, 112 (SAC, 2009-2011) Adequate, 120 (SAC, 2017)
Portugal	PT	None	Adequate, 106 (SAC, 2010)
Slovenia	SI	Mandatory fortification of salt, 19 ppm since 1999	Adequate, 140 (A, 2003-2005)
Spain	ES	Voluntary fortification of salt, 60 ppm since 1983	Adequate, 109 (SAC, 1995, 2000-2002) Adequate, 117 (2009-2010) Adequate, 173 (SAC, 2011-2012)
Sweden	SE	Voluntary fortification of salt, 10 ppm since 1936 50 ppm since 1966	Adequate, 125 (SAC, 2006-2007)
Switzerland	CH	Voluntary fortification of salt, 25 ppm since 2014	Adequate, 115 (SAC, 1999) Adequate, 120 (2009) Adequate, 137 (SAC, 2015)

Abbreviations: UIC urinary iodine concentration, SAC school aged children, A adults, N/A not available

Table 2: Data requested and provided with stratification-level and area.

Country	Country code	Thyroid medication	Thyroid surgery	Radio-iodine	Fine needle biopsies	Diagnoses of thyroid disease	Diagnoses of thyroid cancer
Belgium	BE	Sex/age	Sex/age	Sex/age	Sex/age	N/A	Sex/age
Bulgaria	BG	Sex	Sex	Total	Sex	Sex	Sex/age
Croatia	HR	N/A	N/A	N/A	N/A	N/A	Sex
Czech Republic	CZ	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex
Denmark	DK	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Finland	FI	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Germany	DE	Sex/age*	N/A	N/A	N/A	Sex/age*	N/A
United Kingdom	UK						
<i>Northern Ireland</i>		N/A	N/A	N/A	N/A	N/A	Sex/age
<i>Scotland</i>		N/A	N/A	N/A	N/A	N/A	Sex/age
Greece	GR	Total*	N/A	Total*	N/A	Total*	Total*
Hungary	HU	Sex/age	Sex/age	Sex/age	N/A	Age	Age
Iceland	IS	Sex	N/A	N/A	N/A	N/A	Sex
Ireland	IE	N/A	N/A	N/A	N/A	N/A	Sex/age
Israel	IL	Sex/age	N/A	N/A	N/A	Sex/age	Sex/age
Latvia	LV	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
North Macedonia	MK	Total*	N/A	N/A	N/A	N/A	N/A
Norway	NO	Sex/age	Sex/age	N/A	N/A	Sex/age	Sex/age
Poland	PL	Total*	Total	N/A	N/A	Total	Sex/age
Portugal	PT	Total*	Sex/age	Sex/age*	N/A	Sex/age	Sex/age
Slovenia	SI	Total	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Spain	ES	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Sweden	SE	Sex/age	Sex/age	Total	N/A	Sex/age	Sex/age
Switzerland	CH	N/A	N/A	N/A	N/A	N/A	N/A

 National data

 Regional data or data from well-defined sub-populations

*Data not suitable for inclusion in the database and calculation of raw rates.

Abbreviation: N/A not available

Table 3: Thyroid medication use per 10,000 persons, sex- and age standardized

Country	Country code	Thyroid medication			
		Antithyroid medication, users	Antithyroid medication, DDD	Thyroid hormone therapy, users	Thyroid hormone therapy, DDD
Belgium	BE	19.28		481.46	
Czech Republic	CZ	26.60		612.83	
Denmark	DK	43.19	8,825	241.76	58,356
Finland	FI	9.94		513.95	
Hungary	HU	34.78	4,386	253.09	39,040
Latvia	LV	41.51	5,372	192.36	30,901
Norway	NO	13.12		424.78	
Spain	ES	13.71	1,672	308.24	55,989
Sweden	SE	8.59	3,235	432.84	91,269

Abbreviation: DDD defined daily dose

Table 4: Thyroid surgeries per 10,000 persons, sex- and age standardized

Country	Country code	Thyroid surgery			
		All indications, number of persons	All indications, number of procedures	Benign indications, number of persons	Benign indications, number of procedures
Belgium	BE	5.67		5.05	
Czech Republic	CZ	5.94	6.11	1.42	1.55
Denmark	DK	3.77	4.10	3.37	3.51
Finland	FI	4.12	4.38	3.68	3.79
Hungary	HU	4.57	5.27	4.29	4.83
Latvia	LV	5.20	5.31	4.73	4.76
Norway	NO	3.20	3.28	1.99	2.00
Slovenia	SI			1.57	1.57
Spain	ES	2.77		2.27	
Sweden	SE	5.35	5.53	2.20	2.22

Table 5: Incident thyroid cancer per 10,000 persons, sex- and age standardized

Country	Country code	Diagnoses of thyroid cancer
Belgium	BE	0.87
Bulgaria	BG	0.40
Denmark	DK	1.38
Finland	FI	0.90
Great Britain		
<i>Northern Ireland</i>		0.46
<i>Scotland</i>		0.49
Hungary	HU	0.91
Ireland	IE	0.65
Latvia	LV	1.12
Norway	NO	0.73
Slovenia	SI	0.72
Spain	ES	0.89
Sweden	SE	0.62

Supplementary materials for the paper:

Register-based information on thyroid disease in Europe: lessons and results from the EUthyroid collaboration. By Line Tang Møllehave et al.

Table of contents

Supplementary Table S1: Overview of data provided and first critical assessment

Supplementary Table S2: Overview of data included in the database and second critical assessment

Supplementary Table S3: Raw rates of thyroid medication

Supplementary Table S4: Raw rates of thyroid procedures

Supplementary Table S5: Raw rates of thyroid diagnoses

Supplementary Table S1: Overview of data provided and first critical assessment

Country	Country code	Thyroid medication	Thyroid surgery	Radio-iodine	Fine needle biopsies	Diagnoses of thyroid disease	Diagnoses of thyroid cancer
Belgium	BE	Sex/age	Sex/age	Sex/age	Sex/age	N/A	Sex/age
Bulgaria	BG	Sex	Sex	Total	Sex	Sex	Sex/age
Croatia	HR	N/A	N/A	N/A	N/A	N/A	Sex
Czech Republic	CZ	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex
Denmark	DK	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Finland	FI	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Germany	DE	Sex/age ¹	N/A	N/A	N/A	Sex/age ¹	N/A
United Kingdom	UK						
<i>Northern Ireland</i>		N/A	N/A	N/A	N/A	N/A	Sex/age
<i>Scotland</i>		N/A	N/A	N/A	N/A	N/A	Sex/age
Greece	GR	Total ²	N/A	Total ²	N/A	Total ²	Total ²
Hungary	HU	Sex/age	Sex/age	Sex/age	N/A	Sex/age	Sex/age
Iceland	IS	Sex	N/A	N/A	N/A	N/A	Sex
Ireland	IE	N/A	N/A	N/A	N/A	N/A	Sex/age
Israel	IL	Sex/age	N/A	N/A	N/A	Sex/age	Sex/age
Latvia	LV	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Macedonia	MK	Total ²	N/A	N/A	N/A	N/A	N/A
Norway	NO	Sex/age	Sex/age	N/A	N/A	Sex/age	Sex/age
Poland	PL	Total ²	Total	Total	N/A	Total	Sex/age
Portugal	PT	Total ²	Sex/age	Sex/age	N/A	Sex/age	Sex/age
Slovenia	SI	Total	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Spain	ES	Sex/age	Sex/age	N/A	N/A	Sex/age	Sex/age
Sweden	SE	Sex/age	Sex/age	Total	N/A	Sex/age	Sex/age
Switzerland	CH	N/A	N/A	N/A	N/A	N/A	N/A

N/A not available

Reasons for not included in the database/calculation of raw rates:

¹Cohort-data not related to a specific population, data only related to a small proportion of the total population, or data related to a non-representative sub-population.

²Variables in units that cannot be translated to the requested format.

Supplementary Table S2: Overview of data included in the database and second critical assessment

Country	Country code	Thyroid medication	Thyroid surgery	Radio-iodine	Fine needle biopsies	Diagnoses of thyroid disease	Diagnoses of thyroid cancer
Belgium	BE	Sex/age	Sex/age	Sex/age	Sex/age	N/A	Sex/age
Bulgaria	BG	Sex ¹	Sex ¹	Total ¹	Sex ¹	Sex ¹	Sex/age
Croatia	HR	N/A	N/A	N/A	N/A	N/A	Sex ¹
Czech Republic	CZ	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex ¹
Denmark	DK	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Finland	FI	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
United Kingdom	UK						
<i>Northern Ireland</i>		N/A	N/A	N/A	N/A	N/A	Sex/age
<i>Scotland</i>		N/A	N/A	N/A	N/A	N/A	Sex/age
Hungary	HU	Sex/age	Sex/age	Sex/age	N/A	Sex/age	Sex/age
Iceland	IS	Sex ¹	N/A	N/A	N/A	N/A	Sex ¹
Ireland	IE	N/A	N/A	N/A	N/A	N/A	Sex/age
Israel	IL	Sex/age ³	N/A	N/A	N/A	Sex/age ²	Sex/age ³
Latvia	LV	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Norway	NO	Sex/age	Sex/age	N/A	N/A	Sex/age	Sex/age
Poland	PL	EXC	Total ¹	Total ¹	N/A	Total ¹	Sex/age
Portugal	PT	EXC	Sex/age ²	Sex/age ²	N/A	Sex/age ²	Sex/age ²
Slovenia	SI	Total ¹	Sex/age	Sex/age	Sex/age	Sex/age	Sex/age
Spain	ES	Sex/age	Sex/age	N/A	N/A	Sex/age ²	Sex/age
Sweden	SE	Sex/age	Sex/age	Total ¹	N/A	Sex/age	Sex/age

N/A not available, EXC excluded

Reasons for not included in comparison between countries/calculation of standardized rates:

¹Not sex- and age-stratified

²Based on ICD9-codes

³Invalid datasource or method of registration

Supplementary Table S3: Raw rates of thyroid medication per 10,000 inhabitants

Country	Thyroid medication			
	Antithyroid medication, prevalent users	Antithyroid medication, DDD	Thyroid hormone therapy, prevalent users	Thyroid hormone therapy, DDD
Belgium	19.40		473.51	
Bulgaria	4.79		73.15	
Croatia				
Czech Republic	2.47		659.28	
Denmark	39.71	8146.10	229.17	55615.16
Finland	9.72		508.53	
United Kingdom				
<i>Northern Ireland</i>				
<i>Scotland</i>				
Hungary	33.27	4188.75	251.94	38965.73
Iceland	2.06	208.28	419.79	85166.69
Ireland				
Israel	2.74		99.00	
Latvia	42.17	5463.23	193.72	31087.83
Norway	12.17		379.10	
Poland				
Portugal				
Slovenia		2198.98		52596.04
Spain	13.63	1658.97	293.34	53035.38
Sweden	8.38	3155.68	419.91	88339.69

Supplementary Table S4: Raw rates of thyroid procedures per 10,000 inhabitants

Country	Thyroid surgery				Fine needle biopsies		Radioiodine			
	All indications, number of persons	All indications, number of procedures	Benign indications, number of persons	Benign indications, number of procedures	All indications, number of persons	All indications, number of procedures	All indications, number of persons	All indications, number of doses	Benign indications, number of persons	Benign indications, number of doses
Belgium	5.52		4.91		9.22		0.33		0.30	
Bulgaria	4.07	4.12	3.83	3.75	1.06	1.07	0.44	0.49	0.10	0.18
Croatia										
Czech Republic	6.33	6.51	1.50	1.64	21.47	24.04	1.97		0.28	
Denmark	3.70	4.03	3.31	3.45	1.57	1.80	3.37	3.50	3.23	3.30
Finland	4.10	4.36	3.66	3.77	0.05	0.05	0.62	0.67	0.19	0.20
United Kingdom										
<i>Northern Ireland</i>										
<i>Scotland</i>										
Hungary	4.65	5.36	4.37	4.91			0.51	0.63	0.21	0.25
Iceland										
Ireland										
Israel										
Latvia	5.23	5.35	4.76	4.79	6.58	6.90	2.08	2.17	1.08	1.13
Norway	3.03	3.11	1.90	1.91						
Poland		5.91		5.38			4.13	4.32	3.47	3.59
Portugal	10.84				0.41					
Slovenia			1.56	1.56	12.20	12.20			4.80	4.80
Spain	2.69		2.20						.	.
Sweden	5.20	5.37	2.14	2.15			0.76	2.26	0.58	1.82

Supplementary Table S5: Raw rates of thyroid diagnoses per 10,000 inhabitants

Country	Thyroid cancer	Congenital iodine-deficiency syndrome	Iodine-deficiency-related thyroid disorders and allied conditions	Subclinical iodine-deficiency hypothyroidism	Other hypothyroidism	Other nontoxic goitre	Thyrotoxicosis/Hyperthyroidism	Thyroiditis	Other thyroid disorders	One or more thyroid related diagnoses
Belgium	0.85									
Bulgaria	0.42	0.11	6.28	0.82	113.56	114.22	56.06	45.94	32.70	
Croatia	1.23									
Czech Republic	0.99	1.12	19.71	7.35	296.53	247.51	62.25	186.68	75.60	702.49
Denmark	1.34	0.00	0.06	0.05	17.96	8.05	12.07	2.00	0.37	36.62
Finland	0.90	0.02	0.04	0.04	29.72	7.92	11.09	1.34	0.20	48.96
United Kingdom										
<i>Northern Ireland</i>	0.42									
<i>Scotland</i>	0.47									
Hungary	0.91	4.15	2.59	1.26	50.50	41.78	21.29	9.48	88.72	182.13
Iceland	0.96									
Ireland	0.60									
Israel	14.28	1.78	.	.	173.56	109.44	36.69	30.00	18.61	.
Latvia	1.14	0.39	3.51	1.88	279.81	314.95	107.64	96.81	22.83	615.48
Norway	0.67	0.01	0.04	0.01	5.39	11.04	9.78	2.18	0.29	28.74
Poland	6.99	0.45	16.62	4.89	251.25	128.38	52.59	29.35	42.32	.
Portugal	4.95	0.09	.	.	15.83	11.75	3.45	1.58	.	.
Slovenia	0.78	.	.	.	8.01	9.54	5.87	11.59	.	.
Spain	0.87	0.15	29.25	.	86.15	4.52	67.64	66.71	66.33	.
Sweden	0.60	0.01	0.02	0.02	8.67	7.43	11.48	3.18	0.70	30.25